

SPECTRAL ANALYSIS FOR MATRIX HAMILTONIAN OPERATORS

JEREMY L. MARZUOLA AND GIDEON SIMPSON

ABSTRACT. In this work, we study the spectral properties of matrix Hamiltonians generated by linearizing the nonlinear Schrödinger equation about soliton solutions. By a numerically assisted proof, we show that there are no embedded eigenvalues for the three dimensional cubic equation. Though we focus on a proof of the $3d$ cubic problem, this work presents a new algorithm for verifying certain spectral properties needed to study soliton stability.

Source code for verification of our computations, and for further experimentation, are available at http://www.math.toronto.edu/simpson/files/spec_prop_code.tgz.

CONTENTS

1. Introduction	2
1.1. Main Results	3
1.2. Organization of Results	6
2. Properties of the Nonlinear Schrödinger Equation	6
3. Linearization about a Soliton	8
4. Spectral Properties of the Linearized Hamiltonian	9
4.1. A Survey of Results on the Spectrum of $\mathcal{H}$	9
4.2. Generalized Kernel	14
4.3. Natural Orthogonality Conditions	15
5. Bilinear Forms and the Spectral Property	16
5.1. The Index of an Operator	17
5.2. Numerical Estimates of the Index	18
5.3. Invertibility of Operators	21
5.4. Estimates of Inner Products	22
5.5. Proof of the Spectral Property	22
6. Other Problems	26
6.1. Numerical Estimates of the Index	27
6.2. Estimates of the Inner Products	28

7. Discussion	35
Appendix A. Commutator Estimates	36
A.1. Large Eigenvalues	36
A.2. Spherical Harmonics	37
Appendix B. Proof of the Invertibility of the Operators	40
Appendix C. Numerical Methods	44
C.1. Singularities at the Origin	44
C.2. Artificial Boundary Conditions	45
C.3. Computation of the Indexes	52
C.4. Computation of the Inner Products	53
References	56

1. INTRODUCTION

The nonlinear Schrödinger equation (NLS) in $\mathbb{R} \times \mathbb{R}^d$,

$$(1.1) \quad i\psi_t + \Delta\psi + g(|\psi|^2)\psi = 0, \quad \psi(0, \mathbf{x}) = \psi_0(\mathbf{x}),$$

appears in many different contexts. In applications, it appears as a leading order approximation in nonlinear optics, many body quantum systems, and hydrodynamics. It is also intrinsically interesting as a canonical example of the competition between nonlinearity and dispersion.

For appropriate choices of the nonlinearity $g : \mathbb{R} \rightarrow \mathbb{R}$, the equation is known to possess *soliton* solutions, nonlinear bound states satisfying (1.1) with the *ansatz*

$$\psi(t, \mathbf{x}) = e^{i\lambda t} R(\mathbf{x}; \lambda),$$

where $\lambda > 0$ is the *soliton* parameter. It is conjectured that any solution of (1.1) with appropriate nonlinearity that does not disperse as $t \rightarrow \infty$ must eventually converge to a finite sum of stable solitons. This is referred to as the “soliton resolution” conjecture, a notoriously difficult problem to formulate, see [36].

A natural property to investigate is the stability of the solitons. In [14, 39, 40], a criterion for *orbital* stability is established. Briefly, it says that if the derivative of the L^2 norm of $R(\mathbf{x}; \lambda)$ taken with respect to λ is positive, then the soliton is orbitally stable. The perturbation remains small in a particular norm, H^1 in the case of NLS. If this derivative is negative, the soliton is unstable.

Though these results on the orbital stability of solitons are very powerful, relying on much of the variational structure of the equations, they have three weaknesses. The first is that they do not say if a perturbed soliton reaches an asymptotically constant state;

orbital stability only assures us that the perturbation remains small. The second is that this approach provides no information if the derivative of the L^2 norm vanishes. This is the case of the L^2 critical focusing NLS equation, with $g(s) = s^{d/2}$ and for *saturated* nonlinearities which possess *minimal mass* solitons. Finally, the orbital stability fundamentally depends on the underlying equation possessing a known variational structure. Though this is not a valid criticism for (1.1), it is a problem for other equations, such as those studied in [32].

Alternatively, results such as [3,4,8,16,24,25], and many others, prove *asymptotic* stability of a soliton or a collection of solitons; the system converges to specific solitons as $t \rightarrow \infty$, and the rest of the mass disperses. Asymptotic stability is usually proven perturbatively. The leading order behavior of the perturbation to the soliton is governed by the linearized operator. First, linear stability is proven by assessing the spectrum of the linearized operator. Then the nonlinearity is shown to be dominated by the linear flow. The spectrum of the linearized operator of (1.1) with monomial nonlinearity was studied in [6].

Embedded eigenvalues of the linear operator are detrimental to proving the necessary linear estimates. Indeed, they obstruct the needed dispersive estimates, as demonstrated in [9]. Thus, it is standard to make the assumption that there are no eigenvalues embedded in the essential spectrum. It is known that such a condition cannot be proven directly using abstract properties of the linearized operator, but must in fact be directly related to algebraic properties of the soliton itself. In this work we develop an algorithm for studying the spectral properties of the operator appearing when one linearizes (1.1) about a soliton solution. Furthermore, we use this algorithm to prove the absence of *embedded* eigenvalues or resonances for four NLS problems. We collect these results in the following section.

Remark 1.1. *Though we only prove Theorem 2 for a small number of cases, our objective in this work is to present an approach for verifying the spectral hypotheses required for soliton stability theory.*

1.1. Main Results. Our results hinge on a so called *spectral property* based on linearized matrix Schrödinger operators. The specific form, and its motivations, are developed Section 5. In general, this property can be formulated as:

Definition 1.2 (The Generalized Spectral Property). *Let $d \geq 1$. Given $L_{\pm}$ and a skew adjoint operator Λ , consider the two real Schrödinger operators*

$$\mathcal{L}_+ = -\Delta + \mathcal{V}_+, \quad \mathcal{L}_- = -\Delta + \mathcal{V}_-,$$

defined by

$$\begin{aligned} \mathcal{L}_+ f &= \frac{1}{2} [L_+, \Lambda] f = \frac{1}{2} [L_+ \Lambda f - \Lambda L_+ f], \\ \mathcal{L}_- f &= \frac{1}{2} [L_-, \Lambda] f = \frac{1}{2} [L_- \Lambda f - \Lambda L_- f] \end{aligned}$$

and

$$\mathcal{V}_{\pm} = \frac{1}{2} \mathbf{x} \cdot \nabla V_{\pm}.$$

Let the real quadratic form for $\mathbf{z} = (u, v)^T \in H^1 \times H^1$ be

$$\begin{aligned}\mathcal{B}(\mathbf{z}, \mathbf{z}) &= \mathcal{B}_+(u, u) + \mathcal{B}_-(v, v) \\ &= \langle \mathcal{L}_+ u, u \rangle + \langle \mathcal{L}_- v, v \rangle.\end{aligned}$$

The system is said to satisfy a spectral property on the subspace $\mathcal{U} \subseteq H^1 \times H^1$ if there exists a universal constant $\delta_0 > 0$ such that $\forall \mathbf{z} \in \mathcal{U}$,

$$\mathcal{B}(\mathbf{z}, \mathbf{z}) > \delta_0 \int (|\nabla \mathbf{z}|^2 + e^{-|\mathbf{y}|} |\mathbf{z}|^2) d\mathbf{y}.$$

In this work, the skew adjoint operator is

$$\Lambda f \equiv \frac{d}{2} f + \mathbf{x} \cdot \nabla f = \frac{d}{d\lambda} \left[\lambda^{\frac{d}{2}} f(\lambda \mathbf{x}) \right].$$

This has particular significance for the L^2 critical equation, though we employ it in supercritical problems. This (mis)application is discussed in Section 7.

Our results rely on the key observation of G. Perelman [21] that

Theorem 1. *Given the JL operator, arising from the linearization of NLS about a soliton, where*

$$JL = \begin{pmatrix} 0 & L_- \\ -L_+ & 0 \end{pmatrix}$$

assume the $L_{\pm}$ operators satisfies the Spectral Property in the sense of Definition 1.2. Then JL has no embedded eigenvalues on the subspace $\mathcal{U}$.

Proof. Let us assume we have an embedded eigenstate $\mathbf{z}_{\text{em}} = (u_{\text{em}}, v_{\text{em}})^T \in \mathcal{U}$ corresponding to eigenvalue $i\tau_{\text{em}}$, $\tau_{\text{em}} > \lambda_0$. Then,

$$\begin{aligned}L_- v_{\text{em}} &= i\tau_{\text{em}} u_{\text{em}}, \\ L_+ u_{\text{em}} &= -i\tau_{\text{em}} v_{\text{em}}.\end{aligned}$$

Plugging directly into the form,

$$\begin{aligned}\mathcal{B}(\mathbf{z}_{\text{em}}, \mathbf{z}_{\text{em}}) &= \langle \mathcal{L}_+ u_{\text{em}}, u_{\text{em}} \rangle + \langle \mathcal{L}_- v_{\text{em}}, v_{\text{em}} \rangle \\ &= \frac{1}{2} \{ \langle \Lambda u_{\text{em}}, L_+ u_{\text{em}} \rangle + \langle L_+ u_{\text{em}}, \Lambda u_{\text{em}} \rangle \} \\ &\quad + \frac{1}{2} \{ \langle \Lambda v_{\text{em}}, L_- v_{\text{em}} \rangle + \langle L_- v_{\text{em}}, \Lambda v_{\text{em}} \rangle \} \\ &= \frac{1}{2} \{ i\tau_{\text{em}} \langle \Lambda u_{\text{em}}, v_{\text{em}} \rangle - i\tau_{\text{em}} \langle v_{\text{em}}, \Lambda u_{\text{em}} \rangle \} \\ &\quad + \frac{1}{2} \{ -i\tau_{\text{em}} \langle \Lambda v_{\text{em}}, u_{\text{em}} \rangle + i\tau_{\text{em}} \langle u_{\text{em}}, \Lambda v_{\text{em}} \rangle \} \\ &= \frac{i\tau_{\text{em}}}{2} \{ \langle \Lambda u_{\text{em}}, v_{\text{em}} \rangle - \langle v_{\text{em}}, \Lambda u_{\text{em}} \rangle + \langle v_{\text{em}}, \Lambda u_{\text{em}} \rangle - \langle \Lambda u_{\text{em}}, v_{\text{em}} \rangle \} \\ &= 0.\end{aligned}$$

□

We remark that this holds not just for embedded eigenvalues, but for any purely imaginary eigenvalue. Thus, if the spectral property holds, we are assured that there are no imaginary eigenvalues *on the designated subspace*. The subspace $\mathcal{U}$ will be set by our analysis of the spectrum in Section 4.

Definition 1.3. *Separately, we say that a linearized NLS problem satisfies of a spectral condition if it lacks both:*

- *Embedded eigenvalues,*
- *Endpoint resonances.*

Our second theorem, which relies on the first is:

Theorem 2. *The spectral condition holds for the 3d cubic equation, (1.1) with $d = 3$ and $g(s) = s$, linearized about the ground state soliton R .*

We adapt the methods of [13] to give a numerically assisted proof of this result.

Remark 1.4. *Though the main result of this paper will be to establish Theorem 2, our algorithm can also be used to establish the spectral condition for the one dimensional equation with $g(s) = s^{2.5}$ and $g(s) = s^3$. Again, this is for the problem linearized about the ground state soliton.*

Separately, we establish that for 3d problems with nonlinearities satisfying the conditions necessary for the existence of a soliton, as discussed in [2], one need only test for embedded eigenvalues that are:

- Near the endpoints of the essential spectrum,
- On a sufficiently low spherical harmonic.

In Appendix A we give a proof of the following result using positive commutator arguments otherwise known as Mourre estimates:

Theorem 3. *Given a Hamiltonian $\mathcal{H}$, there exists some $M > 0$ such that for $|\mu| > M$, there are no solutions u_μ such that*

$$\mathcal{H}u_\mu = \mu u_\mu.$$

Similarly, if $d \geq 2$, for any embedded eigenvalue, u_μ , there exists a $K > 0$ such that the spherical harmonic decomposition

$$u_\mu = \sum_{k=0}^{\infty} \alpha_k(r, \mu) \phi_k(\phi, \theta)$$

consists only of harmonics with $k < K$.

Remark 1.5. *Such results are well-known using resolvent estimate techniques; however, our approach provides easily computable limits on M and K in terms of the soliton solution.*

1.2. Organization of Results. In Sections 2, 3 and 4, we review fundamental properties of (1.1) and the associated linearized operator.

In Sections 4.1.2 and 4.1.3, we collect results and adapt the techniques of [12] to prove properties of the discrete spectrum and the absence of embedded resonances.

Finally, in Section 5, we prove an appropriate spectral property as in Definition 1.2, based on the work in [13]. G. Perelman's observation then rules out embedded eigenvalues. This proves Theorem 2.

In Appendices A.1 and A.2, we use Mourre multipliers to eliminate large embedded eigenvalues and large spherical harmonics from the expansion of an embedded eigenvalue. Though these results have been known via resolvent methods for some time, we aim to collect as much analytic information about the spectrum as possible, providing bounds for future estimates and computations. An overview of our numerical methods with benchmarks is then presented in Appendix C.

ACKNOWLEDGMENTS. This paper is an extension of a result of a thesis done by the first author under the direction of Daniel Tataru at the University of California, Berkeley that arose from a discussion with Wilhelm Schlag and Galina Perelman at the Mathematisches Forschungsinstitut Oberwolfach. The first author is supported by an NSF Postdoctoral Fellowship. The second author is supported in part by NSERC. In addition, the authors wish to thank Gadi Fibich, Michael Weinstein, Ian Zwiers and especially Wilhelm Schlag for many helpful conversations throughout the development of the paper.

2. PROPERTIES OF THE NONLINEAR SCHRÖDINGER EQUATION

In this section we briefly review some important properties of (1.1). For additional details, we refer the reader to the texts [5, 35].

In general, for nonlinearity $g : \mathbb{R} \rightarrow \mathbb{R}$, (1.1) possesses the following invariants for data $\psi_0 \in H^1$ and $|\mathbf{x}|\psi_0 \in L^2$:

Conservation of Mass (or Charge):

$$Q(\psi) = \frac{1}{2} \int_{\mathbb{R}^d} |\psi|^2 d\mathbf{x} = \frac{1}{2} \int_{\mathbb{R}^d} |\psi_0|^2 d\mathbf{x},$$

Conservation of Energy:

$$E(\psi) = \int_{\mathbb{R}^d} |\nabla \psi|^2 d\mathbf{x} - \int_{\mathbb{R}^d} G(|\psi|^2) d\mathbf{x} = \int_{\mathbb{R}^d} |\nabla \psi_0|^2 d\mathbf{x} - \int_{\mathbb{R}^d} G(|\psi_0|^2) d\mathbf{x},$$

where

$$G(t) = \int_0^t g(s) ds.$$

Pseudo-Conformal Conservation Law:

$$\|(\mathbf{x} + 2it\nabla)\psi\|_{L^2}^2 - 4t^2 \int_{\mathbb{R}^d} G(|u|^2) d\mathbf{x} = \|\mathbb{X}\psi\|_{L^2}^2 - \int_0^t \theta(s) ds,$$

where

$$\theta(s) = \int_{\mathbb{R}^d} (4(d+2)G(|\psi|^2) - 4dg(|\psi|^2)|\psi|^2) d\mathbf{x}.$$

Note that $(\mathbf{x} + 2it\nabla)$ is the Hamilton flow of the linear Schrödinger equation, so the above identity relates how the solution to the nonlinear equation is effected by the linear flow.

Detailed proofs of these conservation laws can be arrived at easily using energy estimates or Noether's Theorem, which relates conservation laws to symmetries of an equation.

In this work, we restrict our attention to *focusing* nonlinearities, such that $g(s) \geq 0$ for all $s \in \mathbb{R}$. These are the nonlinearities that can yield soliton solutions. Often, $g(s) = s^\sigma$ for some $\sigma > 0$. We examine one instance of the power nonlinearity, the three dimensional cubic problem ($\sigma = 1$).

As noted, a soliton solution takes the form

$$\psi(t, \mathbf{x}) = e^{i\lambda t} R(\mathbf{x}; \lambda),$$

where $\lambda > 0$ and $R(\mathbf{x}; \lambda)$ is a positive, radially symmetric, exponentially decaying solution of the equation:

$$(2.1) \quad \Delta R - \lambda R + g(|R|^2)R = 0.$$

For power nonlinearities, the existence and uniqueness of the ground state soliton is well known. Additionally, the scaling properties of this case permit us to take $\lambda = 1$. We do this in all that follows.

Existence of the soliton is proved by in [2] by minimizing the functional

$$T(\psi) = \int |\nabla \psi|^2 d\mathbf{x}$$

with respect to the constraint of fixed

$$V(\psi) = \int [G(|\psi|^2) - \frac{\lambda}{2} |\psi|^2] d\mathbf{x}.$$

Then, using the minimizing sequence and Schwarz symmetrization, one obtains the existence of the nonnegative, spherically symmetric, decreasing soliton solution. Uniqueness is established in [19] by ODE methods.

An important relation is that $Q(\lambda) = Q(R(\cdot; \lambda))$ and $E(\lambda) = E(R(\cdot; \lambda))$ are differentiable with respect to λ . This fact can be determined from the early works of Shatah, namely [29], [30]. By differentiating Equation (2.1), Q and E with respect to λ , we have

$$\partial_\lambda E = -\lambda \partial_\lambda Q.$$

Variational techniques developed in [14] and [31] tell us that when $\delta(\lambda) = E(\lambda) + \lambda Q(\lambda)$ is convex, or $\delta''(\lambda) > 0$, the soliton is orbitally stable. For $\delta''(\lambda) < 0$ the soliton is unstable to small perturbations. This stability (instability) directly is closely related to the eigenvalues of the matrix Hamiltonians resulting from linearizing NLS about a soliton. For a brief reference on this subject, see [34], Chapter 4.

3. LINEARIZATION ABOUT A SOLITON

Let us write down the form of NLS linearized about a soliton solution. First, we assume we have a solution $\psi = e^{i\lambda t}(R + \phi(\mathbf{x}, t))$. Inserting this into the equation, we have

$$(3.1) \quad i(\phi)_t + \Delta(\phi) = -g(R^2)\phi - 2g'(R^2)R^2\text{Re}(\phi) + O(\phi^2),$$

by splitting ϕ up into its real and imaginary parts, then doing a Taylor Expansion. Hence, if $\phi = u + iv$, we get

$$(3.2) \quad \partial_t \begin{pmatrix} u \\ v \end{pmatrix} = JL \begin{pmatrix} u \\ v \end{pmatrix},$$

where

$$(3.3) \quad JL = \begin{pmatrix} 0 & L_- \\ -L_+ & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} L_+ & 0 \\ 0 & L_- \end{pmatrix}$$

and

$$(3.4) \quad L_- = -\Delta + \lambda - V_-, \quad V_- = g(R),$$

$$(3.5) \quad L_+ = -\Delta + \lambda - V_+, \quad V_+ = g(R) + 2g'(R^2)R^2.$$

Alternatively, if we formulate the problem in terms of ϕ and ϕ^* ,

$$(3.6) \quad \partial_t \begin{pmatrix} \phi \\ \phi^* \end{pmatrix} = i\mathcal{H} \begin{pmatrix} \phi \\ \phi^* \end{pmatrix},$$

where

$$(3.7) \quad \mathcal{H} = \begin{pmatrix} -\Delta + \lambda - V_1 & -V_2 \\ V_2 & \Delta - \lambda + V_1 \end{pmatrix}$$

and

$$(3.8) \quad V_1 = g(R^2) + g'(R^2)R^2, \quad V_2 = g'(R^2)R^2.$$

The potentials in the two formulations are related by $V_+ = V_1 + V_2$ and $V_- = V_1 - V_2$.

There are many things we can immediately say about L_- , L_+ , JL and $\mathcal{H}$. For a reference on the spectral theory involved, see Hislop-Sigal [15] or Reed-Simon [23]. First of all, both L_- and L_+ are self-adjoint operators. Also, L_- is a non-negative definite operator and its null space is $\text{span}\{R\}$. Note also that the functions $\frac{\partial R}{\partial x_j}$ for $j = 1, 2, \dots, d$ are in the null space of L_+ . By comparison with the operator $\Delta + \lambda$ and using the fact that R decays exponentially, we see that the essential spectrum of $\mathcal{H}$ is the set $(-\infty, \lambda] \cup [\lambda, \infty)$.

from Weyl's Theorem, see [12] and [23]. Equivalently, the essential spectrum of JL is $(-i\infty, i\lambda] \cup [i\lambda, i\infty)$. Indeed, $\sigma(JL) = i\sigma(\mathcal{H})$. Finally, using the fact that L_- is non-negative definite and looking at eigenvalues $\mathcal{H}^2$, we see

$$(3.9) \quad L_- L_+ u = \nu^2 u.$$

However, this can be rewritten as

$$Tv = L_-^{\frac{1}{2}} L_+ L_-^{\frac{1}{2}} v = \nu^2 v$$

for $v = L_-^{\frac{1}{2}} u$. Since the operator T is self-adjoint, we must have $\nu \in \mathbb{R} \cup i\mathbb{R}$.

Typically, asymptotic stability is studied with the following assumptions made on the matrix Hamiltonian:

Definition 3.1. *A Hamiltonian, $\mathcal{H}$ is called admissible if the following hold:*

- (1) *There are no embedded eigenvalues in the essential spectrum.*
- (2) *The only real eigenvalue in $[-\lambda, \lambda]$ is 0.*
- (3) *The values $\pm\lambda$ are not resonances.*

Definition 3.2. *Let NLS have nonlinearity g . We call g admissible at λ if there exists a soliton, R_λ , for NLS and the Hamiltonian, $\mathcal{H}$, resulting from linearization about R_λ is admissible in terms of Definition 3.1.*

Remark 3.3. *For simplicity in exposition, a matrix Hamiltonian, $\mathcal{H}$, is said to be admissible if it satisfies several spectral conditions. One of these properties is a lack of embedded eigenvalues in the essential spectrum, which to establish we employ a spectral property as in [13].*

The spectral conditions of $\mathcal{H}$ from Definition 3.1 are generally required to prove dispersive estimates for the evolution operator associated with the linearized Hamiltonian equation, which in turn are required to prove asymptotic stability results for solitons. See [24, 25] for further discussion. Let P_d and P_c be the projections onto the discrete and continuous spectrum of $\mathcal{H}$.

See Figure 1 for a description of the spectral decomposition for $\mathcal{H}$ resulting from linearizing about solitons with subcritical, critical and supercritical stability properties.

4. SPECTRAL PROPERTIES OF THE LINEARIZED HAMILTONIAN

We now give more detailed and formal statements on the spectral properties of the operator under investigation.

4.1. A Survey of Results on the Spectrum of $\mathcal{H}$.

Orbitally Unstable Spectral Decomposition - Exponential Instability (Supercritical Behavior)




FIGURE 1. Plots of the spectral decomposition for $\mathcal{H}$ when R_λ is exponentially unstable (supercritical behavior).

4.1.1. *An Analytic Result on the Spectrum of $\mathcal{H}$.* We formalize the heuristic discussion from Section 3 with the following theorem from [12]. Let us write the operator as

$$\mathcal{H} = \mathcal{H}_0 + V = \begin{bmatrix} -\Delta + \lambda & 0 \\ 0 & \Delta + \lambda \end{bmatrix} + \begin{bmatrix} -V_1 & -V_2 \\ V_2 & V_1 \end{bmatrix}.$$

In [12], the authors proved the following properties of the spectrum:

Theorem 4 (Erdogan-Schlag). *Assume there are no embedded eigenvalues in the continuous spectrum of $\sigma(\mathcal{H})$. The essential spectrum of $\mathcal{H}$ equals $(-\infty, -\lambda] \cup [\lambda, \infty)$. Moreover, $\sigma(\mathcal{H}) = -\sigma(\mathcal{H}) = \overline{\sigma(\mathcal{H})} = \sigma(\mathcal{H}^*)$ and $\sigma(\mathcal{H}) \subset \mathbb{R} \cup i\mathbb{R}$. The discrete spectrum consists of eigenvalues $\{z_j\}_{j=1}^N$, $0 \leq N \leq \infty$, of finite multiplicity. For each $z_j \neq 0$, the algebraic and geometric multiplicities coincide and $\text{Ran}(\mathcal{H} - z_j)$ is closed. The zero eigenvalue has finite multiplicity.*

4.1.2. *Absence of Embedded Resonances.* This result is developed in the earlier work of Erdogan-Schlag [12] and Agmon [1].

Define the space

$$X_\sigma = L^{2,\sigma} \times L^{2,\sigma},$$

where

$$L^{2,\sigma} = \{f \mid |\mathbf{x}|^\sigma f \in L^2\}.$$

Then, we have the following Theorem, proved in [12]:

Theorem 5 (Erdogan-Schlag). *Let V_1, V_2 have sufficient decay at ∞ . Then for any μ such that $|\mu| > \lambda$, $(\mathcal{H}_0 - (\mu \pm i0))^{-1}V : X_{-\frac{1}{2}-} \rightarrow X_{-\frac{1}{2}-}$ is a compact operator, and*

$$I + (\mathcal{H}_0 - (\mu \pm i0))^{-1}V$$

is invertible on these spaces.

The proof relies on a similar argument to a restriction theorem from harmonic analysis, which follows from a calculation using the specific structure of $R_0(z) = (-\Delta - z)^{-1}$. This strategy emulates closely that of the bootstrapping argument of Agmon, [1], for scalar operators.

4.1.3. Discrete Spectrum. We wish to show that the spectrum of $\mathcal{H}$ when linearized about the ground state has the discrete spectral decomposition Figure 1.

Remark 4.1. *For the 3d cubic nonlinearity, the structure of the discrete spectrum away from the essential spectrum has been verified numerically in [10], whose methods we recall briefly here.*

In [25], using arguments derived from [22], it is shown that the discrete spectrum for supercritical exponents is determined by the discrete spectrum of $L_{\pm}$. We present here a slightly stronger version that works for linearizations about a minimal mass soliton, $R = R_{\min}$, in saturated nonlinearities. Though we will not numerically analyze any saturated nonlinearities in the current work, we present the generalized argument saturated nonlinearities are also of interest.

The verification of the discrete spectrum heavily relies on the following result: following

Theorem 6 (Schlag). *Assume that L_- has no discrete eigenvalues on the interval $(0, \lambda]$ and $\mathcal{H}$ is a Hamiltonian as in (3.7) resulting from linearizing about a minimal mass soliton. Then, the only discrete eigenvalue for $\mathcal{H}$ in the interval $[-\lambda, \lambda]$ is 0.*

Remark 4.2. *A very similar theorem appeared in [25] proving the same result for the Hamiltonian $\mathcal{H}$ of the form (3.7) formed from $g(s) = s$ with $\lambda = 1$ and $\mathbf{x} \in \mathbb{R}^3$. The proof follows with minimal changes and is adaptable to many cases, hence we include it below for completeness.*

Proof. We argue by contradiction. To this end, assume $\mathcal{H}$ has an eigenvalue away from 0, say at E . Let $\lambda = 1$ for simplicity. Then $\mathcal{H}^2$ has an eigenvalue at some value $E^2 \in (0, 1]$. Hence, we have

$$L_- L_+ u_E = E^2 u_E,$$

for $E^2 < 1$. Since L_- is self-adjoint, we see that $u_E \perp \phi$. By elliptic regularity, we have that $u_E \in H_{loc}^4$. Let P be the projection orthogonal to ϕ . Let $A = PL_+P$. Using that

$$\ker(L_+) = \text{span}\{\partial_j \phi | 1 \leq j \leq d\}$$

and

$$(4.1) \quad (\mathcal{H} - z)^{-1} = (\mathcal{H}_0 - z)^{-1} [I - U_1 [I - U_2 J(\mathcal{H}_0 - z)^{-1} U_1]^{-1} \\ \times U_2 J(\mathcal{H}_0 - z)^{-1} U_1]^{-1},$$

we have

$$\ker(L_+) = \text{span}\{\partial_\lambda R, \partial_j R | 1 \leq j \leq d\}$$

since

$$\langle R, \partial_\lambda R \rangle = 0.$$

Take E_0 to be the unique negative eigenvalue for L_+ . Then, define

$$g(\alpha) = \langle (L_+ - \alpha)^{-1} R, R \rangle,$$

which is well-defined and differentiable on $(E_0, 1)$ since ϕ is orthogonal to the kernel of L_+ . We have

$$g'(\alpha) = \langle (L_+ - \alpha)^{-1} R, R \rangle > 0$$

and

$$g(0) = \frac{1}{2} \langle R, \partial_\lambda R \rangle = 0.$$

Hence, $g(0) = 0$ is the only 0 for g in the interval $(E_0, 1)$ since

$$\lim_{\alpha \rightarrow E_0} g(\alpha) \rightarrow -\infty.$$

Conversely, if $Af = \alpha f$ for some $-\infty < \alpha < 1$, $\alpha \neq 0$ and $f \in L^2$, then $f \perp R$ and

$$(PL_+P - \lambda)f = (A - \lambda)f = 0.$$

Since

$$E_0 \langle f, f \rangle \leq \langle L_+ f, f \rangle = \lambda \langle f, f \rangle,$$

we know that $\lambda \geq E_0$. If $\lambda = E_0$, then f is a ground state of L_+ and hence not orthogonal to R . However, $g(\lambda) = 0$, hence $\lambda = 0$. So, A has a collection of eigenvalues at 0. Define

$$\mathcal{G} = \text{span}\{R, R_j, R_\lambda, u_E\}.$$

We would like to show that $\dim(\mathcal{G}) = d + 3$. Since ϕ is orthogonal to all the other functions, we need only show that the equation

$$(4.2) \quad c_1 u_E + c_2 R_\lambda + \sum_{j=1}^d c_{j+2} R_j = 0$$

has only the trivial solution $c_j = 0$ for all j . By applying L_+ to (4.2), we see

$$c_1 L_+ u_E + c_2 R = 0.$$

Taking the inner product with u_E , we conclude $c_1 = 0$. This implies that $c_2 = 0$. As a result, $c_j = 0$ for $j = 2, \dots, d+2$. Now, if we can show that

$$(4.3) \quad \sup_{\|f\|_{L^2}, f \in \mathcal{G}} \langle Af, f \rangle < 1,$$

then by the Courant minimax principle, there would be at least $d+3$ eigenvalues less than 1 for A . However, we have shown there are exactly $d+2$ of them. Note that neither the minimal eigenfunction for L_+ nor ϕ itself are eigenvalues of A due to orthogonality arguments. Hence, if we can prove (4.3), we have proved the result.

By our assumption on the spectrum of L_- ,

$$\langle PL_-^{-1}Pf, f \rangle < \langle f, f \rangle$$

for $f \neq 0$. Since $E \leq 1$ by assumption, we can prove the stronger result that

$$\langle Af, f \rangle \leq E^2 \langle PL_-^{-1}Pf, f \rangle$$

for all $f = au_E + b\phi + \mathbf{c} \cdot \nabla R + dR_\lambda$. To this end, we have

$$\begin{aligned} \langle Af, f \rangle &= \langle L_+(au_E), au_E + \mathbf{c} \cdot \nabla \phi + d\phi_\lambda \rangle \\ &\quad + \langle L_+\mathbf{c} \cdot \nabla R, au_E + \mathbf{c} \cdot \nabla R + dR_\lambda \rangle \\ &= E^2 \langle L_-^{-1}(au_E), au_E + \mathbf{c} \cdot \nabla R + dR_\lambda \rangle \\ &\quad + E^2 \langle dR_\lambda, L_-^{-1}(au_E) \rangle \\ &\leq E^2 \langle L_-^{-1}(au_E), au_E + \mathbf{c} \cdot \nabla R + dR_\lambda \rangle + E^2 \langle dR_\lambda, L_-^{-1}(au_E) \rangle \\ &\quad + E^2 \langle L_-^{-1}(\mathbf{c} \cdot \nabla R + dR_\lambda), (\mathbf{c} \cdot \nabla R + dR_\lambda) \rangle \\ &\leq E^2 \langle PL_-^{-1}Pf, f \rangle \end{aligned}$$

since $L_-^{-1}u_E \perp \nabla R$ and L_- is positive definite on $\mathcal{G} \setminus R$. $\square$

In order to test the discrete spectral assumptions, we briefly recall the work of [10], which requires some numerical computation. First, let review the Birman-Schwinger method. let $H = L_- - \lambda = -\Delta - V$ for $V > 0$. Since we are looking for small, positive eigenvalues of L_- , so take $Hf = -\alpha^2 f$ for $0 < \alpha < \lambda$ so we have $L_-f = (\lambda - \alpha^2)f$. Set $U = \sqrt{V}$ and $g = Uf$, then

$$g = U(-\Delta + \alpha^2)^{-1}Ug.$$

In other words, $g \in L^2$ is an eigenfunction for

$$K(\alpha) = U(-\Delta + \alpha^2)^{-1}Ug$$

with eigenvalue 1 where $K > 0$, compact. Conversely, if $g \in L^2$ satisfies $K(\alpha)g = g$, then

$$f = U^{-1}g = (-\Delta + \alpha^2)^{-1}Ug \in L^2$$

and $Hf = -\alpha^2 f$. The eigenvalues of $K(\alpha)$ are seen to be strictly increasing as $\alpha \rightarrow 0$ since

$$K'(\alpha) = -2\lambda U(-\Delta + \alpha^2)^{-2}U.$$

This implies that

$$\# \{ \alpha : \text{Ker}(H - \alpha^2) \neq \{0\} \} = \# \{ E > 1 : \text{Ker}(K(0) - E) \neq \{0\} \}$$

counted with multiplicity.

Finally, use the symmetric resolvent identity to see

$$\begin{aligned} (H - z)^{-1} &= (-\Delta - z)^{-1} + (-\Delta - z)^{-1}U \\ &\quad \times [I - U(-\Delta - z)^{-1}U]^{-1}U(-\Delta - z)^{-1}. \end{aligned}$$

Hence, the Laurent expansion about $z = 0$ does not require negative powers for z iff $I + U(-\Delta - z)^{-1}U$ is invertible at $z = 0$, i.e.

$$\ker(I - U(-\Delta)^{-1}U) = \{0\}$$

by the Fredholm alternative since V has exponential decay. Hence, if H has no resonance or eigenvalue at 0, then $K(0)$ will not have an eigenvalue at 1. If we then count the eigenvalues α_j in decreasing order for $K(0)$, then H has exactly N negative eigenvalues and neither an eigenvalue or resonance at 0 iff $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_N > 1$ and $\alpha_{N+1} < 1$. Hence, we can study numerically study K for a soliton of the saturated nonlinear Schrödinger equation. To do so, we must accurately find a soliton, then use it as potential in the truncation and discretization scheme presented in [10], where the gap condition is verified for the Hamiltonian resulting from linearization about the 3d cubic ground state soliton.

4.2. Generalized Kernel. Let us review the generalized kernel of a Hamiltonian resulting from linearizing about a soliton. Following [39], we see by direct calculation that the vectors

$$\begin{bmatrix} 0 \\ R \end{bmatrix}, \begin{bmatrix} R_j \\ 0 \end{bmatrix}$$

for all $j = 1, \dots, d$ are contained in $\ker(JL)$. Now, as $Q(R_\lambda)$ is differentiable with respect to λ , we have by a simple calculation that $L_+ \partial_\lambda R = -R$ and $L_-(x\phi) = -2\nabla R$. Hence, the vectors

$$\begin{bmatrix} 0 \\ x_j R \end{bmatrix}, \begin{bmatrix} (\partial_\lambda R)_{\lambda_0} \\ 0 \end{bmatrix}$$

in the generalized null space of order 2. Notice that so far we have constructed at $2d + 2$ dimensional null space. Since we know the null spaces of L_- and L_+ exactly, these are unique. For power nonlinearities, $g(s) = s^\sigma$,

$$(4.4) \quad (\partial_\lambda R)_{\lambda_0=1} = \frac{1}{2} \left(\frac{1}{\sigma} R + \mathbf{x} \cdot \nabla R \right).$$

We use this explicit form in our calculations.

As a result, we have the following

Theorem 7. *Let g be the L^2 supercritical monomial nonlinearity. There exists a $2d + 2$ dimensional null space for $\mathcal{H}$, the matrix Hamiltonian resulting from linearization about the ground state soliton ($\lambda = 1$), consisting of the span of the vectors*

$$\left\{ \begin{bmatrix} 0 \\ R \end{bmatrix}, \begin{bmatrix} R_j \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ x_j R \end{bmatrix}, \begin{bmatrix} (\partial_\lambda R)_{\lambda_0} \\ 0 \end{bmatrix} \right\}.$$

Proof. The generalized null space of the adjoint can be found by reversing the location of the non-zero elements in the above vectors.

Suppose that there exists a generalized eigenspace for the eigenvalue $E \neq 0$. Then, there exists $\chi \neq 0$ and $\psi \neq 0$ such that $(\mathcal{H} - E)\psi = \chi$ and $(\mathcal{H} - E)\chi = 0$. Then, note that

$$\begin{aligned} (\mathcal{H}^2 - E^2)\psi &= (A + E)\chi = 2E\chi, \\ (\mathcal{H}^2 - E^2)\chi &= 0. \end{aligned}$$

Hence, $\mathcal{H}^2$ has a generalized eigenspace at E^2 . As a result, we see that $T = L_+ L_-$ has a generalized eigenspace at E^2 . Let $T\chi = E^2\chi$ and $(T - E^2)\psi = c\chi$, for some $c \neq 0$. Hence,

$$\begin{aligned} (L_-^{\frac{1}{2}} L_+ L_-^{\frac{1}{2}} - E^2) L_-^{\frac{1}{2}} \psi_1 &= c\chi_1, \\ (L_-^{\frac{1}{2}} L_+ L_-^{\frac{1}{2}} - E^2)^2 L_-^{\frac{1}{2}} \psi_1 &= c L_-^{\frac{1}{2}} (L_+ L_- - E^2) \chi_1 = 0, \end{aligned}$$

where given $P_R^c = I - P_R$, we have $\chi_1 = P_R^c \chi \neq 0$ since $T\chi = E^2\chi$ and $\psi_1 = P_R^c \psi \neq 0$ since $(T - E^2)\psi = c\chi$. However, this means that the self-adjoint operator $L_-^{\frac{1}{2}} L_+ L_-^{\frac{1}{2}}$ has a generalized eigenvalue, which is impossible by an orthogonality argument.

Since we have assumed there are no eigenvalues at the endpoints of the continuous spectrum, there can be no accumulation and the number of discrete eigenvalues is finite. $\square$

4.3. Natural Orthogonality Conditions. As noted in Definition 1.2, even if the spectral property holds, it only implies the absence of embedded eigenvalues on a subspace of $\mathcal{U} \subset L^2 \times L^2$. That we must limit ourselves to a subspace will become clear in Section 5.2, where we demonstrate that the operators $\mathcal{L}_\pm$ have negative eigenvalues.

This subspace will be defined as the orthogonal complement to the span of a set of vectors. If this collection of vectors is not chosen properly, we may find that the spectral property holds though the operator still has embedded eigenvalues. Thus the constraints on the set of vectors whose orthogonal complement will define $\mathcal{U}$ are:

- (1) They must be orthogonal to any embedded eigenvalues,
- (2) Orthogonality with respect to them should induce positivity of $\mathcal{L}$ on $\mathcal{U}$.

A way of meeting both of these requirements is to use the discrete spectrum of the adjoint matrix Hamiltonian, $\mathcal{H}^*$. To that end, we rely on the following simple results.

Lemma 4.3. *If $(\lambda, \vec{u})$ is an eigenvalue, eigenvector pair for JL and $(\sigma, \vec{v})$ is an eigenvalue, eigenvector pair for $(JL)^*$, then*

$$(\lambda - \sigma^*) \langle \vec{u}, \vec{v} \rangle = 0.$$

Thus, if $\lambda - \sigma^* \neq 0$, the states are orthogonal.

Corollary 4.4. *An eigenstate of JL associated with an imaginary (possibly embedded) eigenvalue, $i\tau \neq 0$, is orthogonal to $\ker_g((JL)^*)$.*

Corollary 4.5. *Let $(i\tau \neq 0, \vec{\psi})$ and $(\lambda > 0, \vec{\phi})$ be eigenvalue, eigenvector pairs of JL . Then*

$$\begin{aligned} \langle \psi_1, \phi_2 \rangle &= 0, \\ \langle \psi_2, \phi_1 \rangle &= 0. \end{aligned}$$

Proof. By the Hamiltonian symmetry of the problem, $-\lambda (\phi_2, \phi_1)^T$ and $\lambda, (-\phi_2, -\phi_1)^T$ are eigenvalue pairs of the adjoint, $(JL)^*$. Therefore,

$$\begin{aligned} \langle \psi_1, \phi_2 \rangle - \langle \psi_2, \phi_1 \rangle &= 0, \\ -\langle \psi_1, \phi_2 \rangle - \langle \psi_2, \phi_1 \rangle &= 0. \end{aligned}$$

Adding and subtracting these equations gives the result. $\square$

These trivial observations motivate using the known spectrum of the adjoint system in constructing the orthogonal subspace. For the 3D cubic problem, we can thus use make use of eigenstates coming from the origin and the two off axis, real eigenvalues.

5. BILINEAR FORMS AND THE SPECTRAL PROPERTY

We show here how the Spectral Property 1.2 is a condition sufficient for showing there are no embedded eigenvalues, thus proving Theorems 1 and 2.

For general nonlinearities, we assume the operator resulting from linearizing about a soliton R has as discrete spectrum that is one of the following:

- (i) a $2d + 2$ dimensional null space given by $R, \nabla R, \partial_\lambda R, \mathbf{x}R$ plus 2 eigenfunctions with symmetric discrete eigenvalues $\lambda_0, -\lambda_0$ such that $\lambda_0^2 \in \mathbb{R}$,
- (ii) a $2d + 4$ dimensional null space given by $R, \nabla R, \partial_\lambda R, \mathbf{x}R, \alpha, \beta$.

In the following subsections, we establish the following:

Theorem 8. *The generalized spectral property holds for the 3d cubic problem for $(f, g)^T \in \mathcal{U} \subset L^2 \times L^2$ specified by the following orthogonality conditions:*

$$\langle f, R \rangle = 0, \quad \langle g, R + \mathbf{x} \cdot \nabla R \rangle = 0, \quad \langle f, \phi_2 \rangle = 0, \quad \langle f, x_j R \rangle = 0 \quad \text{for } j = 1, \dots, d,$$

where $\vec{\phi} = (\phi_1, \phi_2)^T$ is the eigenstate associated with the positive eigenvalue $\sigma > 0$.

This subspace is motivated by the observations in Section 4.3, as all of the elements we are orthogonal to arise from the spectrum of the adjoint problem.

Remark 5.1. *Note that henceforward we assume here the unstable eigenfunction for the 3d cubic problem is radial. This claim is substantiated by direct integration in our numerical results section, as well as the fact that the spectral decomposition for the critical problem remains valid under the assumption of radial symmetry, hence the multiplicity of radial eigenfunctions must be at least 4. Since R , R_λ are the only radial components of the kernel, the unstable eigenmode must also be radial.*

Following [13], we computationally verify the spectral property in the following steps. First, the bilinear form is decomposed by spherical harmonics into

$$(5.1) \quad \begin{aligned} \mathcal{B}(\mathbf{z}, \mathbf{z}) &= \mathcal{B}_+(f, f) + \mathcal{B}_-(g, g) \\ &= \sum_{k=0}^{\infty} \mathcal{B}_+^{(k)}(f^{(k)}, f^{(k)}) + \sum_{k=0}^{\infty} \mathcal{B}_-^{(k)}(g^{(k)}, g^{(k)}), \end{aligned}$$

where $\mathbf{z} = (f, g)^T$, and $f^{(k)}$ and $g^{(k)}$ are the components of f and g in the k -th spherical harmonic. We then identify the dimension of the subspace of negative eigenvalues for $\mathcal{L}_\pm^{(k)}$. Though at first this would appear to require an infinite number of computations, a monotonicity property of these operators with respect to k limits this to a finite number of harmonics. We then show that our orthogonality conditions are sufficient to point us away from the negative directions, allowing us to prove our result.

5.1. The Index of an Operator. For a bilinear form B on a vector space V , the index of B with respect to V is given by

$$\text{ind}_V(B) \equiv \max\{k \in \mathbb{N} \mid \text{there exists a subspace } P \text{ of codimension } k \text{ such that } B|_P \text{ is positive}\}.$$

Our results rely on the following generalization of Theorem XIII.8 of [23], which is in turn an extension of the Sturm Oscillation Theorem (Section XIII.7 of [23]):

Theorem 9. *Let $U^{(k)}$ be the solution to*

$$L^{(k)}U^{(k)} = -\frac{d^2}{dr^2}U^{(k)} - \frac{d-1}{r}\frac{d}{dr}U^{(k)} + V(r)U^{(k)} + \frac{k(k+d-1)}{r^2}U^{(k)} = 0$$

with initial conditions given by the limits

$$\lim_{r \rightarrow 0} \frac{U^{(k)}(r)}{r^k} = 1, \quad \lim_{r \rightarrow 0} \frac{d}{dr} \frac{U^{(k)}(r)}{r^k} = 0,$$

where V is sufficiently smooth and decaying at ∞ . Then, the number $N(U^{(k)})$ of zeros of $U^{(k)}$ is finite and

$$\begin{aligned} \text{ind}_{H_{\text{rad}}^1}(B^{(0)}) &= N(U^{(0)}), \\ \text{ind}_{H_{\text{rad}+}^1}(B^{(k)}) &= N(U^{(k)}), \quad k \geq 1, \end{aligned}$$

where $B^{(k)}$ is the bilinear form associated to $L^{(k)}$.

The space H_{rad}^1 is the set of radially symmetric $H^1(\mathbb{R}^d)$ functions. The space $H_{\text{rad}+}^1$ is the subset of H_{rad}^1 for which

$$\int \frac{|f|^2}{|\mathbf{x}|^2} d\mathbf{x} < \infty.$$

We will omit the subscript notation in our subsequent index computations. It will be H_{rad}^1 for $k = 0$ and $H_{\text{rad}+}^1$ for $k \geq 1$.

If one wishes to remove the limits from the statement of the initial conditions, let $U^{(k)}(r) = r^k \tilde{U}^{(k)}(r)$. Then the operator becomes

$$\tilde{L}^{(k)} = -\frac{d^2}{dr^2} - \frac{d-1+2k}{r} \frac{d}{dr} + V$$

and the initial conditions become $\tilde{U}^{(k)}(0) = 1$ and $\frac{d}{dr} \tilde{U}^{(k)}(0) = 0$. Indeed, we use precisely this change of variables when making our numerical computations; see Appendix C.1. The proof can be adapted from the proof of Theorem XIII.8 in [23].

Corollary 5.2. *The index is monotonic with respect to k ,*

$$\text{ind}(B^{(k+1)}) \leq \text{ind}(B^{(k)}).$$

This has the useful consequence that once we find an k for which $\text{ind}(B^{(k)}) = 0$, we can immediately conclude that $B^{(k')} \geq 0$ for all $k' \geq k$.

Once we have computed the number of directions of each $\mathcal{L}_{\pm}^{(k)}$ that prevents it from being positive, we can check that we have a sufficient number of orthogonal conditions to point us into the positive subspace.

5.2. Numerical Estimates of the Index. To compute the indexes of the operators, we proceed as follows.

In dimension three, we solve the initial value problems

$$(5.2) \quad \mathcal{L}_+^{(0)} U^{(0)} = 0, \quad U^{(0)}(0) = 1, \quad \frac{d}{dr} U^{(0)}(0) = 0,$$

$$(5.3) \quad \mathcal{L}_-^{(0)} Z^{(0)} = 0, \quad Z^{(0)}(0) = 1, \quad \frac{d}{dr} Z^{(0)}(0) = 0$$

for radially symmetric functions $U^{(0)}$ and $Z^{(0)}$. For higher harmonics, $k > 0$, we solve the initial value problems

$$(5.4) \quad \mathcal{L}_+^{(k)} U^{(k)} = 0, \quad U^{(k)}(0) = 0, \quad \lim_{r \rightarrow 0} r^{-k} U^{(k)}(r) = 1,$$

$$(5.5) \quad \mathcal{L}_-^{(k)} Z^{(k)} = 0, \quad Z^{(k)}(0) = 0, \quad \lim_{r \rightarrow 0} r^{-k} Z^{(k)}(r) = 1$$

for radially symmetric functions $U^{(k)}$ and $Z^{(k)}$.

Proposition 5.3. *The indexes of 3d Cubic NLS are:*

$$\begin{aligned} \text{ind}\mathcal{L}_+^{(0)} &= 1, & \text{ind}\mathcal{L}_+^{(1)} &= 1, & \text{ind}\mathcal{L}_+^{(2)} &= 0, \\ \text{ind}\mathcal{L}_-^{(0)} &= 1, & \text{ind}\mathcal{L}_-^{(1)} &= 0. \end{aligned}$$

Once this proposition is established, Corollary 5.2 immediately gives us

Corollary 5.4. *For 3d Cubic NLS,*

$$\begin{aligned} \text{ind}\mathcal{B}_+^{(k)} &= 0 \quad \text{for } k > 2, \\ \text{ind}\mathcal{B}_-^{(k)} &= 0 \quad \text{for } k > 1, \end{aligned}$$

where $\mathcal{B}_\pm^{(k)}$ is the bilinear form associated with $\mathcal{L}_\pm^{(k)}$.

Using the method discussed in Section C, we compute $U^{(\alpha)}$ and $Z^{(\alpha)}$ for each problem, $\alpha = 0, 1, 2$. The profiles appear in Figures 2,3,4. All were computed with a tolerance setting 10^{-13} .

As a consistency check on the numerics, we note that asymptotically, the potential vanishes, and

$$(5.6) \quad \mathcal{L}_\pm^{(k)} \approx -\frac{d^2}{dr^2} - \frac{d-1}{r}U^{(k)} + \frac{k(k+d-2)}{r^2}.$$

In the region where $r \gg 1$, and the equations are essentially free and the solutions must behave as:

$$(5.7a) \quad U^{(k)}(r) \approx C_0^{(k)}r^k + C_1^{(k)}r^{2-d-k}$$

and

$$(5.7b) \quad Z^{(k)}(r) \approx D_0^{(k)}r^k + D_1^{(k)}r^{2-d-k}.$$

Estimating these constants from the numerics, we see that they have the “correct” signs. For instance in Figure 2 (a), $U^{(0)}$ clearly has one zero crossing. For sufficiently large r , the function appears to be increasing past a local minimum. However, since the constants appear to have stabilized, we contend we have entered the free region; the signs and magnitudes of the constants thus forbid another zero.

Proposition 5.5. *For the operators in Proposition 5.3 and Corollary 5.4 there exists a universal $\delta_0 > 0$, sufficiently small, such that for the perturbed operators*

$$\overline{\mathcal{L}}_\pm^{(k)} = \mathcal{L}_\pm^{(k)} - \delta_0 e^{-|\mathbf{x}|^2}$$

the associated bilinear forms have the property that

$$\text{ind}(\overline{\mathcal{B}}_\pm^{(k)}) = \text{ind}(\mathcal{B}_\pm^{(k)}).$$




FIGURE 2. Index computations for 3d Cubic NLS. The number of zero crossings (other than $r = 0$), determines the codimension of the subspace on which the operator $\mathcal{L}_{\pm}^{(k)}$ is positive.




FIGURE 3. Index computations for 3d Cubic NLS. The number of zero crossings (other than $r = 0$), determines the codimension of the subspace on which the operator $\mathcal{L}_{\pm}^{(k)}$ is positive.

Proof. The proof here follows obviously from definition of the index of B , namely the positivity of the quadratic form B on the subspace for individual operators. Let δ_0 be a sufficiently small value that it holds for $\overline{\mathcal{L}}_+^{(k)}$ for $k = 0, 1, 2$ and $\overline{\mathcal{L}}_-^{(k)}$ for $k = 0, 1$. This δ_0 now holds for all higher values of k , again by a monotonicity argument. Indeed, for $k > 2$,

$$\overline{\mathcal{B}}_+^{(k)}(f, f) = \overline{\mathcal{B}}_+^{(2)}(f, f) + \int \frac{k^2 + 2k - 8}{|\mathbf{x}|^2} |f|^2 d\mathbf{x} \geq 0.$$




FIGURE 4. Index computations for 3d Cubic NLS. The number of zero crossings (other than $r = 0$), determines the codimension of the subspace on which the operator $\mathcal{L}_{\pm}^{(k)}$ is positive.

Because the form remains positive, this confirms that its index of zero is unperturbed. $\square$

5.3. Invertibility of Operators. In conjunction with the results on the indexes of operators, we need to compute a number of inner products of the form $\langle \mathcal{L}u, u \rangle$, where $\mathcal{L}$ is one of our operators and u solves $\mathcal{L}u = f$. These are computed numerically, but we can rigorously justify the existence and uniqueness of these solutions, u , for the problems under consideration.

Proposition 5.6 (Numerically Verified for 3d Problems). *Let f be a smooth, radially symmetric, localized function satisfying the bound $|f(r)| \leq Ce^{-\kappa r}$ for some positive constants C and κ . There exists a unique radially symmetric solution*

$$(1 + r^{k+1})u \in L^\infty([0, \infty)) \cap C^2([0, \infty))$$

to

$$(5.8) \quad \mathcal{L}u = f,$$

where $\mathcal{L} = \mathcal{L}_{\pm}^{(k)}$ for one of the 3d problems.

Proof. This is Proposition 2 and 4 of [13], along with our computations of the indexes in Proposition 5.3. See Appendix B for a proof in 1d. $\square$

Corollary 5.7. *The solutions in Proposition 5.6 are smooth and decay $\propto r^{-1-k}$ as $r \rightarrow \infty$.*

5.4. Estimates of Inner Products. In order to prove the spectral property for each of these NLS equations, we need to approximate the bilinear forms associated with $\mathcal{L}_\pm^{(\alpha)}$ on certain functions. These particular functions are, generically, of the form $\mathcal{L}u = f$, where f is from one of the orthogonality conditions.

Proposition 5.8 (Numerical approximation of inner products). *For the 3d cubic problem, let $U_1^{(0)}$, $U_2^{(0)}$, $U_1^{(1)}$, and $Z_1^{(0)}$ solve*

$$(5.9) \quad \mathcal{L}_+^{(0)} U_1^{(0)} = R, \quad (1+r)U_1^{(0)} \in L^\infty,$$

$$(5.10) \quad \mathcal{L}_+^{(0)} U_2^{(0)} = \phi_2, \quad (1+r)U_2^{(0)} \in L^\infty,$$

$$(5.11) \quad \mathcal{L}_+^{(1)} U_1^{(1)} = rR, \quad (1+r^2)U_1^{(0)} \in L^\infty,$$

$$(5.12) \quad \mathcal{L}_-^{(0)} Z_1^{(0)} = R + rR', \quad (1+r)Z_1^{(0)} \in L^\infty,$$

where $\vec{\phi}$ is the eigenstate associated with the positive real eigenvalue of JL . Then,

$$(5.13) \quad K_1^{(0)} \equiv \langle \mathcal{L}_+^{(0)} U_1^{(0)}, U_1^{(0)} \rangle = 1.04846,$$

$$(5.14) \quad K_2^{(0)} \equiv \langle \mathcal{L}_+^{(0)} U_2^{(0)}, U_2^{(0)} \rangle = 0.00215981,$$

$$(5.15) \quad K_3^{(0)} \equiv \langle \mathcal{L}_+^{(0)} U_1^{(0)}, U_2^{(0)} \rangle = -0.116369,$$

$$(5.16) \quad K_1^{(1)} \equiv \langle \mathcal{L}_+^{(1)} U_1^{(1)}, U_1^{(1)} \rangle = -0.581854,$$

$$(5.17) \quad J_1^{(0)} \equiv \langle \mathcal{L}_-^{(0)} Z_1^{(0)}, Z_1^{(0)} \rangle = -0.662038.$$

Proof. These result follows from direct computation. $\square$

Finally, we state the following

Proposition 5.9. *For each case in Proposition 5.8, there exists a δ_0 sufficiently small such that inner products associated with $\bar{\mathcal{L}}$ can be made arbitrarily close to $\mathcal{L}$. These values will be denoted with overlines.*

Proof. This follows immediately from the invertibility of the operator and continuity. $\square$

5.5. Proof of the Spectral Property. We are now ready to prove the spectral property. We prove positivity of $\bar{\mathcal{B}}_+^{(0)}$, the other cases are similar. Our proof closely follows Step 1 and Step 3 of Section 2.4 of [13].

Since $K_1^{(0)}$ and $K_2^{(0)} > 0$, orthogonality to R and ϕ_2 will not give positivity. However, if f is orthogonal to both of these, then it is also orthogonal to

$$q = R - \frac{K_3^{(0)}}{K_2^{(0)}} \phi_2$$

and

$$\begin{aligned}
\langle \mathcal{L}_+^{(0)} q, q \rangle &= K_1^{(0)} - 2 \frac{K_3^{(0)}}{K_2^{(0)}} K_3^{(0)} + \left(\frac{K_3^{(0)}}{K_2^{(0)}} \right)^2 K_2^{(0)} \\
&= -\frac{1}{K_2^{(0)}} ((K_3^{(0)})^2 - K_1^{(0)} K_2^{(0)}) \\
&= -5.22138.
\end{aligned}$$

By Proposition 5.9, we can take δ_0 sufficiently small such that

$$-\frac{1}{\overline{K}_2^{(0)}} \left((\overline{K}_3^{(0)})^2 - \overline{K}_1^{(0)} \overline{K}_2^{(0)} \right) < 0.$$

We proceed with this value of δ_0 .

Let $\overline{Q}$ solve

$$\overline{\mathcal{L}}_+^{(0)} \overline{Q} = q.$$

Obviously,

$$Q = \overline{U}_1^{(0)} - \frac{K_3^{(0)}}{K_2^{(0)}} \overline{U}_2^{(0)}$$

and

$$\overline{\mathcal{B}}_+^{(0)}(\overline{Q}, \overline{Q}) < 0.$$

For a moment, suppose $\overline{Q} \in H_{\text{rad}}^1$; it is not since it decays too slowly to be in L^2 . We could then imagine decomposing H_{rad}^1 into $\text{span}\{\overline{Q}\}$ and its orthogonal complement, where the orthogonalization is done with respect to the $\overline{\mathcal{B}}_+^{(0)}$ quadratic form. Since $\overline{\mathcal{B}}_+^{(0)}(\overline{Q}, \overline{Q}) < 0$, the form is *non-degenerate* and this decomposition is well defined. Since $\text{ind}_{H_{\text{rad}}^1} \overline{\mathcal{B}}_+^{(0)} = 1$, $\overline{\mathcal{B}}_+^{(0)} \geq 0$ on $\text{span}\{\overline{Q}_A\}^\perp$. To prove this claim, we argue by contradiction. Suppose there were an element, $Z \in \text{span}\{\overline{Q}_A\}^\perp$ for which $\overline{\mathcal{B}}_+^{(0)}(Z, Z) < 0$. Then, because of our decomposition, $\overline{\mathcal{B}}_+^{(0)} < 0$ on $\text{span}\{Z, \overline{Q}_A\}$, a space of dimension 2. This contradicts our index calculation, proving the claim.

Continuing, if $u \in H_{\text{rad}}^1$, $u \perp q$ (with respect to L^2), then using the hypothetical orthogonal decomposition,

$$u = c\overline{Q} + u^\perp.$$

If $c = 0$, then u lies in a subspace of H_{rad}^1 on which $\overline{\mathcal{B}}_+^{(0)} \geq 0$, giving the desired positivity. Indeed, the orthogonality condition, $u \perp q$, is sufficient to ensure u is orthogonal to $\overline{Q}$ with respect to the $\overline{\mathcal{B}}_+^{(0)}$ quadratic form. Taking the inner product of u with q ,

$$\begin{aligned}
0 &= c \langle \overline{Q}, q \rangle + \langle u^\perp, q \rangle = c \overline{\mathcal{B}}_+^{(0)}(\overline{Q}, \overline{Q}) + \langle u^\perp, \overline{\mathcal{L}}_+^{(0)} \overline{Q} \rangle \\
&= c \overline{\mathcal{B}}_+^{(0)}(\overline{Q}, \overline{Q}) + \overline{\mathcal{B}}_+^{(0)}(u^\perp, \overline{Q}) = c \overline{\mathcal{B}}_+^{(0)}(\overline{Q}, \overline{Q}) + 0.
\end{aligned}$$

Since $\bar{\mathcal{B}}_+^{(0)}(\bar{Q}, \bar{Q}) \neq 0$, we have $c = 0$.

Unfortunately, the above argument does not work as stated because $\bar{Q}$ is not in L^2 ! To get positivity of $\bar{\mathcal{B}}_+^{(0)}$, we regularize the problem and follow the above scheme. First, we introduce the smooth cutoff function $\chi_A(r) = \chi(r/A)$, defined such that

$$\chi(r) = \begin{cases} 1 & r < 1 \\ 0 & r \geq 2 \end{cases}$$

and the norm

$$\|f\|_{\pm}^2 = \|\nabla f\|_{L^2}^2 + \int |\mathcal{V}_{\pm}| |f|^2.$$

Let $\bar{Q}_A(r) = \bar{Q}(r)\chi_A(r)$. Next, we observe that

$$(5.18) \quad \lim_{A \rightarrow +\infty} \|\bar{Q}_A - \bar{Q}\|_+ + |\bar{\mathcal{B}}_+^{(0)}(\bar{Q}, \bar{Q}) - \bar{\mathcal{B}}_+^{(0)}(\bar{Q}_A, \bar{Q}_A)| = 0.$$

Since $\bar{\mathcal{B}}_+^{(0)}(\bar{Q}, \bar{Q}) < 0$, for sufficiently large A , $\bar{\mathcal{B}}_+^{(0)}(\bar{Q}_A, \bar{Q}_A) < 0$ too. Thus, we can legitimately decompose H_{rad}^1 as

$$(5.19) \quad H_{\text{rad}}^1 = \text{span}\{\bar{Q}_A\} \oplus \text{span}\{\bar{Q}_A\}^{\perp}$$

with the orthogonalization is done with respect to the quadratic form $\bar{\mathcal{B}}_+^{(0)}$.

Finally, let $u \in H_{\text{rad}}^1$, $u \perp R$ and $u \perp \phi_2$. Then $u \perp q$. With A sufficiently large to make the above decomposition valid,

$$(5.20) \quad u = c(A)\bar{Q}_A + u_A^{\perp}.$$

As in the heuristic argument $\bar{\mathcal{B}}_+^{(0)}(u_A^{\perp}, u_A^{\perp}) \geq 0$. Thus,

$$\bar{\mathcal{B}}_+^{(0)}(u, u) = c(A)^2 \bar{\mathcal{B}}_+^{(0)}(\bar{Q}_A, \bar{Q}_A) + \bar{\mathcal{B}}_+^{(0)}(u_A^{\perp}, u_A^{\perp}) \geq c(A)^2 \bar{\mathcal{B}}_+^{(0)}(\bar{Q}_A, \bar{Q}_A).$$

We will have our result if $c(A) \rightarrow 0$ as $A \rightarrow +\infty$.

Since $\langle u, q \rangle = 0$,

$$\begin{aligned} c(A) \langle \bar{U}_A^*, q \rangle &= -\langle u_A^{\perp}, q \rangle \\ &= -\langle u_A^{\perp}, \bar{\mathcal{L}}_+^{(0)} \bar{Q} \rangle \\ &= -\langle u_A^{\perp}, \bar{\mathcal{L}}_+^{(0)} (\bar{Q} - \bar{Q}_A) \rangle. \end{aligned}$$

Therefore,

$$|c(A)| = \frac{|\langle u_A^{\perp}, \bar{\mathcal{L}}_+^{(0)} (\bar{Q} - \bar{Q}_A) \rangle|}{|\langle \bar{U}_A^*, q \rangle|} \leq \frac{\|u_A^{\perp}\|_+ \|\bar{Q} - \bar{Q}_A\|_+}{|\langle \bar{U}_A^*, q \rangle|}.$$

Also,

$$|\langle \bar{U}_A^*, q \rangle - \langle \bar{Q}, q \rangle| = |\langle \bar{U}_A^* - \bar{Q}, \bar{\mathcal{L}}_+^{(0)} \bar{Q} \rangle| \leq \|\bar{Q} - \bar{Q}_A\|_+ \|\bar{Q}\|_+.$$

Because this vanishes as $A \rightarrow +\infty$, we have that for all A sufficiently large,

$$|c(A)| \leq C \|u_A^\perp\|_+ \|\bar{Q} - \bar{Q}_A\|_+$$

for a constant C independent of A .

By construction,

$$\|u_A^\perp\|_+ \leq C \left(\|u\|_+ + c(A) \|\bar{U}_A^*\|_+ \right) \leq C \left(\|u\|_+ + c(A) \|\bar{U}_A^* - \bar{Q}\|_+ + \|\bar{Q}\|_+ \right).$$

Substituting into our previous estimate on $c(A)$,

$$|c(A)| \leq C \left(\|u\|_+ + c(A) \|\bar{U}_A^* - \bar{Q}\|_+ + \|\bar{Q}\|_+ \right) \|\bar{Q} - \bar{Q}_A\|_+.$$

We can clearly see that as $A \rightarrow +\infty$, $c(A) \rightarrow +0$. We conclude,

$$\bar{\mathcal{B}}_+^{(0)}(u, u) \geq 0$$

for $u \in H_{\text{rad}}^1$ and $u \perp R$ and $u \perp \phi_2$. This yields the estimate

$$\mathcal{B}_+^{(0)}(u, u) \geq \delta_0 \int e^{-|\mathbf{x}|^2} |u|^2 dx.$$

Following the same analysis for $\bar{\mathcal{L}}_-$, we conclude

$$\mathcal{B}_-^{(0)}(g, g) \geq \delta_0 \int e^{-|\mathbf{x}|^2} |g|^2 dx$$

for $g \in H_{\text{rad}}^1$ and $g \perp R + rR'$ since $J_1^{(0)} < 0$. Repeating this again for $f \in H_{\text{rad}(1)}^1$, $f \perp rR$, we get

$$\mathcal{B}_-^{(1)}(f, f) \geq \delta_0 \int e^{-|\mathbf{x}|^2} |f|^2 dx$$

because $K_1^{(1)} < 0$.

Let us assume that δ_0 has been taken sufficiently small such that:

- The indexes of all operators are unperturbed, as in Proposition 5.5,
- The above arguments on the positivity of $\bar{\mathcal{B}}_+^{(k)}$ for $k = 0, 1$ and $\bar{\mathcal{B}}_-^{(0)}$ hold.

Then for $\mathbf{z} = (f, g)^T$ satisfying the orthogonality conditions,

$$\begin{aligned}
\overline{\mathcal{B}}(\mathbf{z}, \mathbf{z}) &= \overline{\mathcal{B}}_+(f, f) + \overline{\mathcal{B}}_-(g, g) \\
&= \sum_{k=0}^{\infty} \overline{\mathcal{B}}_+^{(k)}(f^{(k)}, f^{(k)}) + \sum_{k=0}^{\infty} \overline{\mathcal{B}}_-^{(k)}(g^{(k)}, g^{(k)}) \\
&= \sum_{k=0}^{\infty} \mathcal{B}_+^{(k)}(f^{(k)}, f^{(k)}) + \sum_{k=0}^{\infty} \mathcal{B}_-^{(k)}(g^{(k)}, g^{(k)}) - \delta_0 \int e^{-|\mathbf{x}|^2} (|f|^2 + |g|^2) d\mathbf{x} \\
&= \mathcal{B}(\mathbf{z}, \mathbf{z}) - \delta_0 \int e^{-|\mathbf{x}|^2} |\mathbf{z}|^2 d\mathbf{x} \geq 0.
\end{aligned}$$

We almost have the expression in Definition 1.2. To complete the proof, note that for any $\theta \in (0, 1)$,

$$(1 + \theta)\mathcal{B}(\mathbf{z}, \mathbf{z}) \geq \theta \left(\int |\nabla \mathbf{z}|^2 d\mathbf{x} + \int V_+ |f|^2 + \int V_- |g|^2 \right) + \delta_0 \int e^{-|\mathbf{x}|^2} |\mathbf{z}|^2 d\mathbf{x}.$$

We can take $\theta = \theta_*$ sufficiently small such that

$$\theta_* \left(\int V_+ |f|^2 + \int V_- |g|^2 \right) + \delta_0 \int e^{-|\mathbf{x}|^2} |\mathbf{z}|^2 d\mathbf{x} \geq \frac{\delta_0}{2} \int e^{-|\mathbf{x}|^2} |\mathbf{z}|^2 d\mathbf{x}.$$

Then,

$$\mathcal{B}(\mathbf{z}, \mathbf{z}) \geq \frac{\theta_*}{1 + \theta_*} \int |\nabla \mathbf{z}|^2 d\mathbf{x} + \frac{\delta_0}{2(1 + \theta_*)} \int e^{-|\mathbf{x}|^2} |\mathbf{z}|^2 d\mathbf{x}.$$

Shrinking δ_0 again, so that it is smaller than

$$\min \left\{ \frac{\theta_*}{1 + \theta_*}, \frac{\delta_0}{2(1 + \theta_*)} \right\}$$

gives us the spectral property. □

6. OTHER PROBLEMS

In principle, this scheme can be applied to any linearized nonlinear Schrödinger equation. One finds the indexes of the operators, picks an appropriate subspace to project away from, and computes the necessary inner products. However, our experiments show that the algorithm is not as universal as might be hoped. In this section we exhibit the computations for several 1d NLS equations,

$$(6.1) \quad i\psi_t + \psi_{xx} + |\psi|^{2\sigma}\psi = 0.$$

Sometimes our approach works, ruling out embedded eigenvalues in a range of supercritical cases, while in others it fails, leaving a large range of interesting problems unresolved.

6.1. Numerical Estimates of the Index. As in the 3d problem, we first compute the indexes of the operators $\mathcal{L}_\pm$ to identify the number of “bad” directions. In contrast to the multidimensional problems where there are an arbitrarily high, but finite, number of harmonics which must be examined, 1D problems only require us to study the operators restricted to even and odd functions. This requires the following results, whose proofs are quite similar to that of Theorem 9:

Corollary 6.1. *Let U be the even solution to*

$$\begin{cases} LU = -U'' + V(r)U = 0, \\ U(0) = 1, U'(0) = 0, \end{cases}$$

where V is sufficiently smooth and decaying at ∞ . Then, the number $N(U)$ of zeros of U is finite and

$$\text{ind}_{H_e^1}(B) = N(U),$$

where B is the bilinear form associated with L .

Corollary 6.2. *Let U be the odd solution to*

$$\begin{cases} LU = -U'' - \frac{d-1}{r}U' + V(r)U = 0, \\ U(0) = 0, U'(0) = 1, \end{cases}$$

where V is sufficiently smooth and decaying at ∞ . Then, the number $N(U)$ of zeros of U is finite and

$$\text{ind}_{H_o^1}(B) = N(U),$$

where B is the bilinear form associated with L .

H_e^1 and H_o^1 are the subspaces of $H^1(\mathbb{R})$ restricted to even and odd functions. In what follows, we shall omit them in the subscripts of the indexes.

To proceed, we numerically solve the initial value problems

$$(6.2) \quad \mathcal{L}_+^{(e)}U^{(e)} = 0, \quad U^{(e)}(0) = 1, \quad \frac{d}{dx}U^{(e)}(0) = 0,$$

$$(6.3) \quad \mathcal{L}_-^{(e)}Z^{(e)} = 0, \quad Z^{(e)}(0) = 1, \quad \frac{d}{dx}Z^{(e)}(0) = 0$$

for even functions $U^{(e)}$ and $Z^{(e)}$. We then solve

$$(6.4) \quad \mathcal{L}_+^{(o)}U^{(o)} = 0, \quad U^{(o)}(0) = 0, \quad \frac{d}{dx}U^{(o)}(0) = 1,$$

$$(6.5) \quad \mathcal{L}_-^{(o)}Z^{(o)} = 0, \quad Z^{(o)}(0) = 0, \quad \frac{d}{dx}Z^{(o)}(0) = 1$$

for odd functions $U^{(o)}$ and $Z^{(o)}$. $\mathcal{L}_\pm$ have the same definitions as before; the Laplacian is now one dimensional. As in the 3d case, we verify *a posteriori* that, asymptotically, U^α and




FIGURE 5. Index computations for critical 1d NLS. The number of zero crossings (other than $x = 0$), determines the codimension of the subspace on which the operator $\mathcal{L}_{\pm}^{(e/o)}$ is positive.

Z^{α} fit

$$(6.6) \quad U^{\alpha(x)} \approx C_0^{(\alpha)} + C_1^{(\alpha)}x,$$

$$(6.7) \quad Z^{\alpha(x)} \approx D_0^{(\alpha)} + D_1^{(\alpha)}x.$$

In addition, the constants have the right signs to ensure we have the correct number of zero crossings.

Proposition 6.3 (Numerically Verified). *The indexes for the 1d NLS equation with $\sigma = 2, 2.1, 2.5, 3$ are:*

$$\begin{aligned} \text{ind}\mathcal{L}_+^{(e)} &= 1, & \text{ind}\mathcal{L}_-^{(e)} &= 1, \\ \text{ind}\mathcal{L}_+^{(o)} &= 1, & \text{ind}\mathcal{L}_-^{(o)} &= 0. \end{aligned}$$

Proof. Using the method discussed in Section C, we compute $U^{(\alpha)}$ and $Z^{(\alpha)}$ for each problem, $\alpha = e, o$. The profiles appear in Figures 7, 8. All are computed with a relative tolerance of 10^{-10} and an absolute tolerance of 10^{-12} using MATLAB.

□

Proposition 5.5 applies to these 1d problems too. As in the 3d case, we ultimately use a perturbed bilinear form in the proof of the spectral property.

6.2. Estimates of the Inner Products. We now compute a series of inner products and show that in some cases the natural orthogonality conditions are sufficient to yield a spectral




FIGURE 6. Index computations for critical 1d NLS. The number of zero crossings (other than $x = 0$), determines the codimension of the subspace on which the operator $\mathcal{L}_{\pm}^{(e/o)}$ is positive.




FIGURE 7. Index computations for 1d NLS with $\sigma = 2.1$. The number of zero crossings (other than $x = 0$), determines the codimension of the subspace on which the operator $\mathcal{L}_{\pm}^{(e/o)}$ is positive.

property. Rigourously, these results require the following Proposition on the invertibility of the $\mathcal{L}_{\pm}^{(e/o)}$ operators:

Proposition 6.4 (Numerically Verified for 1d Saturated Problem). *Let f be a smooth, localized function satisfying the bound $|f(x)| \leq Ce^{-\kappa|x|}$ for some positive constants C and*




FIGURE 8. Index computations for 1d NLS with $\sigma = 2.1$. The number of zero crossings (other than $x = 0$), determines the codimension of the subspace on which the operator $\mathcal{L}_{\pm}^{(e/o)}$ is positive.




FIGURE 9. Index computations for 1d NLS with $\sigma = 2.5$. The number of zero crossings (other than $x = 0$), determines the codimension of the subspace on which the operator $\mathcal{L}_{\pm}^{(e/o)}$ is positive.

κ . If f is even/odd, there exists a unique even/odd solution $u \in L^\infty(\mathbb{R}) \cap C^2(\mathbb{R})$ to (6.8)

$$\mathcal{L}u = f,$$

where $\mathcal{L} = \mathcal{L}_{\pm}^{(e/o)}$ for a 1d problem.

Proof. See Appendix B. □




FIGURE 10. Index computations for 1d NLS with $\sigma = 2.5$. The number of zero crossings (other than $x = 0$), determines the codimension of the subspace on which the operator $\mathcal{L}_{\pm}^{(e/o)}$ is positive.




FIGURE 11. Index computations for 1d NLS with $\sigma = 3$. The number of zero crossings (other than $x = 0$), determines the codimension of the subspace on which the operator $\mathcal{L}_{\pm}^{(e/o)}$ is positive.




FIGURE 12. Index computations for 1d NLS with $\sigma = 3$. The number of zero crossings (other than $x = 0$), determines the codimension of the subspace on which the operator $\mathcal{L}_{\pm}^{(e/o)}$ is positive.

Proposition 6.5 (Numerical). *Let $U_1^{(e)}$, $Z_1^{(e)}$, and $U_1^{(o)}$, all elements of $L^\infty(\mathbb{R})$, solve the following boundary value problems:*

$$(6.9) \quad \mathcal{L}_+^{(e)} U_1^{(e)} = R, \quad \frac{d}{dx} U_1^{(e)}(0) = 0,$$

$$(6.10) \quad \mathcal{L}_-^{(e)} Z_1^{(e)} = \frac{1}{\sigma} R + x R', \quad \frac{d}{dx} Z_1^{(e)}(0) = 0,$$

$$(6.11) \quad \mathcal{L}_+^{(o)} U_1^{(o)} = R', \quad U_1^{(o)}(0) = 0.$$

Let

$$(6.12) \quad K_1^{(e)} \equiv \mathcal{B}_+^{(e)}(U_1^{(e)}) = \langle \mathcal{L}_+^{(e)} U_1^{(e)}, U_1^{(e)} \rangle,$$

$$(6.13) \quad J_1^{(e)} \equiv \mathcal{B}_-^{(e)}(Z_1^{(e)}) = \langle \mathcal{L}_-^{(e)} Z_1^{(e)}, Z_1^{(e)} \rangle,$$

$$(6.14) \quad K_1^{(o)} \equiv \mathcal{B}_+^{(o)}(U_1^{(o)}) = \langle \mathcal{L}_+^{(o)} U_1^{(o)}, U_1^{(o)} \rangle.$$

Then,

σ	$K_1^{(e)}$	$J_1^{(e)}$	$K_1^{(o)}$
2.0	-0.557768	0.292551	-1.30410
2.1	-0.496932	0.216284	-1.21364
2.5	-0.297841	-0.0216292	-0.924662
3.0	-0.122559	-0.218499	-0.671783.

6.2.1. *Proof of the Spectral Property for Certain Supercritical Cases.* We restrict our attention to the 1d supercritical problems $\sigma = 2.5$ and $\sigma = 3$. Repeating the procedure of Section 5.5, for $\mathbf{z} = (f, g)^T$, the orthogonality of f to R and xR gives us $\bar{\mathcal{L}}_+ \geq 0$ and the orthogonality of g to $\frac{1}{\sigma}R + xR'$ gives us $\bar{\mathcal{L}}_- \geq 0$. This proves the spectral property on the restricted subspace. Since these orthogonality conditions are consistent with those formulated in Section 4.3, we conclude that there are no non zero purely imaginary eigenvalues.

6.2.2. *An Inconclusive Supercritical Case.* In the case of $\sigma = 2.1$, we have that $J_1^{(e)} > 0$, which means that orthogonality of g with respect to $\frac{1}{\sigma}R + xR'$, is insufficient to guarantee positivity of $\mathcal{L}_-$. It is possible that if we extend our scope, as in the 3D cubic problem, to include orthogonality to the eigenstate associated with the unstable eigenvalue we will be able to prove the spectral property for this problem. However, we do not pursue that here; rather we wish to highlight the failure of our algorithm at a seemingly arbitrary supercritical nonlinearity.

6.2.3. *The Critical Case.* The critical 1D problem, with $\sigma = 2$, is also inconclusive. As in the supercritical problems we will look at the inner products against R and $\frac{1}{\sigma}R + xR'$. We also employ inner products arising from with the rest of the generalized kernel, x^2R and β , where β solves

$$L_+\beta = -x^2R.$$

See [39] for details. This motivates the following numerical result:

Proposition 6.6 (Numerical). *Let $Z_2^{(e)}$ solve*

$$(6.15) \quad \mathcal{L}_-^{(e)} Z_2^{(e)} = \rho, \quad \frac{d}{dx} Z_1^{(e)}(0) = 0$$

and let

$$(6.16) \quad J_2^{(e)} \equiv \mathcal{B}_-^{(e)}(Z_2^{(e)}, Z_2^{(e)}) = \left\langle \mathcal{L}_-^{(e)} Z_2^{(e)}, Z_2^{(e)} \right\rangle,$$

$$(6.17) \quad J_3^{(e)} \equiv \mathcal{B}_-^{(e)}(Z_1^{(e)}, Z_2^{(e)}) = \left\langle \mathcal{L}_-^{(e)} Z_1^{(e)}, Z_2^{(e)} \right\rangle.$$

Then,

$$(6.18) \quad J_2^{(e)} = 3.77915,$$

$$(6.19) \quad J_3^{(e)} = 0.864273.$$

Since $K_1^{(e)} < 0$, we may conclude that $\mathcal{L}_+ \geq 0$, when the operator is restricted to even functions that are orthogonal to R . However, orthogonality to neither $\frac{1}{2}R + xR'$ nor ρ is, individually, sufficient to gain positivity of $\mathcal{L}_-^{(e)}$. We are thus motivated to consider

orthogonality to the subspace $\text{span}\{\frac{1}{2}R + xR', \rho\}$, as in the proof of the 3D cubic problem. We examine the quantity

$$(6.20) \quad -\frac{1}{J_2^{(e)}} \left((J_3^{(e)})^2 - J_1^{(e)} J_2^{(e)} \right) = 0.0948958.$$

However, we need this to be negative. Thus, we have no set of natural orthogonality conditions which yield a spectral property. In this case, since $K_1^{(o)} < 0$, the only obstacle to the proof is the $\mathcal{L}_-^{(e)}$ operator.

6.2.4. The Critical Case with Other Orthogonality Conditions. If we had instead used the orthogonality condition, $g \perp R$, and then solved the boundary value problem $\mathcal{L}_-^{(e)} \check{Z}_3^{(e)} = R$, the inner product,

$$\check{J}_1^{(e)} \equiv \left\langle \mathcal{L}_-^{(e)} \check{Z}_1^{(e)}, \check{Z}_1^{(e)} \right\rangle = -3.770731.$$

This would give us a spectral property, but it is not a convenient subspace.

Suppose we use the orthogonality conditions of [13], and let $g \perp \Lambda R$ and $g \perp \Lambda^2 R$. Then, we compute as follows: Let $\hat{Z}_1^{(e)}$, and $\hat{Z}_2^{(e)}$ solve the following problems:

$$(6.21) \quad \mathcal{L}_-^{(e)} \hat{Z}_1^{(e)} = \Lambda R, \quad \frac{d}{dx} \hat{Z}_1^{(e)}(0) = 0,$$

$$(6.22) \quad \mathcal{L}_-^{(e)} \hat{Z}_2^{(e)} = \Lambda R, \quad \frac{d}{dx} \hat{Z}_2^{(e)}(0) = 0,$$

and define the inner products

$$(6.23) \quad \hat{J}_1^{(e)} \equiv \mathcal{B}_-^{(e)}(\hat{Z}_1^{(e)}, \hat{Z}_1^{(e)}) = \left\langle \mathcal{L}_-^{(e)} \hat{Z}_1^{(e)}, \hat{Z}_1^{(e)} \right\rangle,$$

$$(6.24) \quad \hat{J}_2^{(e)} \equiv \mathcal{B}_-^{(e)}(\hat{Z}_2^{(e)}, \hat{Z}_2^{(e)}) = \left\langle \mathcal{L}_-^{(e)} \hat{Z}_2^{(e)}, \hat{Z}_2^{(e)} \right\rangle,$$

$$(6.25) \quad \hat{J}_3^{(e)} \equiv \mathcal{B}_-^{(e)}(\hat{Z}_1^{(e)}, \hat{Z}_2^{(e)}) = \left\langle \mathcal{L}_-^{(e)} \hat{Z}_1^{(e)}, \hat{Z}_2^{(e)} \right\rangle.$$

We will find that

$$(6.26) \quad \hat{J}_1^{(e)} = 0.292551,$$

$$(6.27) \quad \hat{J}_2^{(e)} = 2.57656,$$

$$(6.28) \quad \hat{J}_3^{(e)} = -1.27657.$$

As one would hope, given that the 1D spectral property was established in [20],

$$(6.29) \quad -\frac{1}{\hat{J}_2^{(e)}} \left((\hat{J}_3^{(e)})^2 - \hat{J}_1^{(e)} \hat{J}_2^{(e)} \right) = -0.339932.$$

This sign ensures that projection away from those two directions is sufficient to point us away from the negative eigenvalue, rendering $\overline{\mathcal{L}}_-^{(e)} \geq 0$.

7. DISCUSSION

We have demonstrated a computer assisted algorithm for proving the positivity of a bilinear form, $\mathcal{B}$, on a subspace $\mathcal{U}$. Because of the relationship between $\mathcal{B}$, $\mathcal{L}$, and the linearized operator, JL , we infer that there are no embedded eigenvalues. We succeeded with this program in the case of the 3d cubic equation, and a two supercritical 1d problems. C. Sulem has suggested that it is likely to also be successful for solitons (with $\lambda = 1$) of the 3d cubic-quintic equation,

$$i\psi_t + \Delta\psi + |\psi|^2\psi - \gamma|\psi|^4\psi = 0$$

for γ sufficiently close to zero. We also anticipate success for other 1d supercritical problems with σ sufficiently large. These cases warrant further study.

For subcritical problems, a similar algorithm should apply, though it will certainly require additional orthogonality conditions. Many subcritical problems contain eigenvalues inside the spectral gap. Since our approach does not distinguish between embedded eigenvalues and imaginary eigenvalues in the gap, it would be essential to project away from those states.

It remains to be seen how to extend our technique to other NLS/GP equations. Indeed, the failure in the 1D critical problem is curious. The success or failure of the approach is likely related to the choice of our operator $\Lambda = d/2 + \mathbf{x} \cdot \nabla$. In [13], the authors proved the spectral property using this Λ , as it is generated by the scaling invariance of the mass critical problem. This results in the so-called “pseudoconformal invariant” for critical NLS and has great implications for blow-up. See [20] and [35] for additional details.

Finally, recall that Λ determines the operators $\mathcal{L}_\pm$. These each have an index identifying the number of negative directions. We then choose orthogonality conditions that simultaneously must satisfy the two properties:

- (1) They must be orthogonal to any embedded eigenvalues,
- (2) Orthogonality in L^2 with respect to these directions must imply orthogonality to the negative directions of $\mathcal{L}_\pm$, with respect to the $\mathcal{B}_\pm$ quadratic form.

The first requirement is satisfied by the vectors from the adjoint problem, as discussed in Section 4.3. We appear to have little flexibility in altering these. Changing Λ will change $\mathcal{L}_\pm$; in turn this changes the negative directions. Thus, a different skew adjoint operator may extend the applicability of the algorithm.

APPENDIX A. COMMUTATOR ESTIMATES

A.1. Large Eigenvalues. We now establish upper bounds on the magnitude of embedded L^2 eigenvalues of JL . In this analysis, the fourth order equation comes from squaring the operator. Therefore, we study eigenfunctions $u \in L^2$ of the differential operator:

$$(A.1) \quad L_- L_+ u = \mu^2 u,$$

where the $L_\pm$ operators are

$$L_\pm = -\Delta + \lambda - V_\pm$$

and $\mu \in \sigma_{\text{cont.}}(JL) = (\lambda, \infty)$. As our proof applies to many NLS equations, we do not further specify the potentials $V_\pm$; they are defined as in Section 3.

From the properties of the soliton and the nonlinearity, we have that any solution to equation A.1 is locally smooth via an iteration argument [18]. Following [2], asymptotic analysis shows that a solution decays exponentially fast. As a result, the possible range of frequencies is limited by

$$(A.2) \quad \|\nabla u\|_{L^2} \leq \mu \|u\|_{L^2}.$$

See [33, 37] for references on microlocal analysis. We prove the following:

Theorem 10. *There exists a $\mu_0 > \lambda$ such that for all $\mu \geq \mu_0$, the eigenvalue equation (A.1) has only the trivial zero solution in L^2 .*

Proof. We begin by defining the standard Mourre commutator as:

$$(A.3) \quad M \equiv \mathbf{x} \cdot \nabla + \nabla \cdot \mathbf{x}.$$

Using the structure of these operators, we immediately have the two identities

$$(A.4) \quad \langle [ML_- L_+ + L_+ L_- M]u, u \rangle = 0,$$

$$(A.5) \quad \langle [L_- L_+ - \mu^4]u, u \rangle = 0,$$

where $\langle \cdot, \cdot \rangle$ is the standard L^2 inner product. Combining these two identities with the frequency bound of (A.2), we can rule out L^2 solutions for sufficiently large μ . Indeed,

$$\begin{aligned} \langle (\Delta^2 - 2\lambda\Delta + \Delta(V_+(\mathbf{x})) + V_-(\mathbf{x})\Delta - \lambda(V_- + V_+) \\ + (\lambda^2 - \mu^2) + V_- V_+)u, u \rangle = 0, \quad (*) \end{aligned}$$

$$\begin{aligned} ML_- L_+ - L_+ L_- M &= [M, L_-]L_+ + L_-[M, L_+] + [L_-, L_+]M, \\ [M, -\Delta] &= 4\Delta, \quad [M, V_-] = 2\mathbf{x} \cdot \nabla V_-. \end{aligned}$$

Hence,

$$\begin{aligned} \langle (-8\Delta^2 + 8\lambda\Delta - 4(V_- + V_+)\Delta + 2(\mathbf{x} \cdot \nabla V_- + \mathbf{x} \cdot \nabla V_+)\Delta \\ - 2\lambda(\mathbf{x} \cdot \nabla V_- + \mathbf{x} \cdot \nabla V_+) + 2(\mathbf{x} \cdot \nabla V_-)V_+ + 2(\mathbf{x} \cdot \nabla V_+)V_- \quad (**) \\ - [(V_- - V_+)\Delta - \Delta(V_- - V_+)](d + 2\mathbf{x} \cdot \nabla)u, u \rangle = 0. \end{aligned}$$

Since the last term is a product of skew-adjoint operators, they commute.

Combining 4(*) + (**) and with the frequency bound $\|\nabla u\|_{L^2} \leq \mu^2 \|u\|_{L^2}$, we have for $\mu > \mu_0$, this is a negative definite system. Thus, there are no eigenvalues. Furthermore, this estimate combined and standard Sobolev embeddings implies $u \in H^k$ for any k ; hence, u is smooth.

The system we assess is:

$$\begin{aligned} \int & [-4(\Delta u)^2 - 4\lambda(V_- + V_+)u^2 - 4(\mu^2 - \lambda^2)u^2] \\ & + [4V_-V_+ - 2\lambda(\mathbf{x} \cdot \nabla V_- + \mathbf{x} \cdot \nabla V_+) \\ & + 2(\mathbf{x} \cdot \nabla V_-)V_+ + 2(\mathbf{x} \cdot \nabla V_+)V_- + \Delta(\mathbf{x} \cdot \nabla V_- + \mathbf{x} \cdot \nabla V_+) \\ & - \mathbf{x} \cdot \nabla(\Delta V_- - \Delta V_+) - d(\Delta V_- - \Delta V_+)]u^2 \\ & + [4(\mathbf{x} \cdot \nabla u)(\nabla(V_- - V_+) \cdot \nabla u) \\ & - 2(\mathbf{x} \cdot \nabla V_- + \mathbf{x} \cdot \nabla V_+)\nabla u \cdot \nabla u] d\mathbf{x} = 0. \end{aligned}$$

Hence,

$$\begin{aligned} \int & (-4(\Delta u)^2 - 4\lambda(V_- + V_+)u^2) d\mathbf{x} - 4(\mu^2 - \lambda^2)\|u\|_{L^2}^2 + \|F\|_{L^\infty}\|u\|_{L^2}^2 \\ & + (2\|\mathbf{x} \cdot \nabla V_- + \mathbf{x} \cdot \nabla V_+\|_{L^\infty} \\ & + C_d \max_{j,k} \|\partial_j(V_- - V_+)x_k\|)\|\nabla u\|_{L^2}^2 \\ & \leq \int (-4(\Delta u)^2 - 4\lambda(V_- + V_+)u^2) d\mathbf{x} \\ & - 4(\mu^2 - \lambda^2 - C_1 - C_2\mu^2)\|u\|_{L^2}^2, \end{aligned}$$

where

$$\begin{aligned} F &= 4V_-V_+ - 2\lambda(\mathbf{x} \cdot \nabla V_- + \mathbf{x} \cdot \nabla V_+) + 2(\mathbf{x} \cdot \nabla V_-)V_+ \\ &+ 2(\mathbf{x} \cdot \nabla V_+)V_- + \Delta(\mathbf{x} \cdot \nabla V_- + \mathbf{x} \cdot \nabla V_+) \\ &- \mathbf{x} \cdot \nabla(\Delta V_- - \Delta V_+) - d(\Delta V_- - \Delta V_+) \end{aligned}$$

and $C_j = C_j(V_-, V_+, \lambda, d)$ for $j = 1, 2$. Hence, for μ large, we have that:

$$\int (-4(\Delta u)^2 - 4\lambda(V_- + V_+)u^2 - C_3(\mu, \lambda, V_-, V_+)u^2) d\mathbf{x} \leq 0,$$

for $C_3 > 0$ and u a smooth function, hence $u = 0$. $\square$

A.2. Spherical Harmonics. As our potentials are radially symmetric, we expand our functions in spherical harmonics. Separating the radial variable, r , from the angular variables, θ , the expansion takes the form

$$\sum_k u_k(r) \phi_k(\theta),$$

where

$$\Delta_S \phi_k(\theta) = (k^2 + (d-1)d) \phi_k(\theta).$$

See [38] for a description of the eigenspaces of the spherical Laplacian, Δ_S . Then, we have the following ODE eigenvalue problem:

$$(A.6) \quad \left[\left(-\frac{\partial^2}{\partial r^2} - \frac{d-1}{r} \frac{\partial}{\partial r} + \frac{\alpha^2}{r^2} + \lambda - V_-(r) \right) \times \right. \\ \left. \left(-\frac{\partial^2}{\partial r^2} - \frac{d-1}{r} \frac{\partial}{\partial r} + \frac{\alpha^2}{r^2} + \lambda - V_+(r) \right) - \mu^2 \right] u_k(r) = 0,$$

where $\alpha^2 = k^2 + (d-1)d$ for $k = 0, 1, 2, \dots$. Note, we use the notation α^2 since all of the eigenvalues are non-negative, which will be important in the sequel.

We have following theorem:

Theorem 11. *There exists some $\alpha_0 > 0$ such that for all $\alpha \geq \alpha_0$, the eigenvalue equation (A.6) has only the trivial solution in L^2 .*

Proof. Let us denote the radial inner product by:

$$\langle u, v \rangle_r = \int u v r^{d-1} dr,$$

and the operators:

$$\Delta_r = r^{1-d} \frac{\partial}{\partial r} \left(r^{d-1} \frac{\partial}{\partial r} \right), \\ P_r = d + 2r \frac{\partial}{\partial r}.$$

Using the same commutator approach as in Section A.1,

$$\begin{aligned} & \langle [4\Delta_r - \frac{4\alpha^2}{r^2} - 2rV'_-(r)][-\Delta_r + \frac{\alpha^2}{r^2} + \lambda - V_+(r)] \\ & + [-\Delta_r + \frac{\alpha^2}{r^2} + \lambda - V_-(r)][4\Delta_r - \frac{4\alpha^2}{r^2} - 2rV'_+(r)] \\ & + [(V_- - V_+)\Delta_r - \Delta_r(V_- - V_+)] [d + 2r \frac{\partial}{\partial r}] u_k, u_k \rangle_r = 0. \end{aligned}$$

Thus,

$$\begin{aligned}
\langle -8(\Delta_r)^2 u &+ 16\Delta_r\left(\frac{\alpha^2}{r^2}u_k\right) + 8\lambda\Delta_ru_k - 4(V_- + V_+)\Delta_ru_k - \frac{8\alpha^4}{r^4}u_k \\
&- \frac{8\lambda\alpha^2}{r^2}u_k + \frac{4\alpha^2}{r^2}(V_- + V_+)u_k + 2(rV'_-(r) + rV'_+(r))\Delta_ru_k \\
&- 2\frac{\alpha^2}{r}(V'_-(r) + V'_+(r))u_k - 2\lambda r(V'_-(r) + V'_+(r))u_k \\
&+ 2rV'_-(r)V_+(r)u_k + 2rV_-(r)V'_+(r)u_k \\
&+ (d + 2r\frac{\partial}{\partial r})((V_- - V_+)\Delta_r - \Delta_r(V_- - V_+))u_k, u_k \rangle_r \\
&= 0.
\end{aligned}$$

This implies:

$$\begin{aligned}
\int [-8(\Delta_ru_k)^2 &- (16\frac{\alpha^2}{r^2} + 8\lambda)((u_k)_r)^2 - (\frac{8\alpha^4}{r^4} + \frac{8\lambda\alpha^2}{r^2})u_k^2]r^{d-1}dr \\
&+ \int [4(V_- + V_+) + 2r(V_-)_r - 6r(V_+)_r]((u_k)_r)^2r^{d-1}dr \\
&+ \int [\frac{4\alpha^2}{r^2}(V_- + V_+) - (d-4)\frac{\alpha^2}{r^4} - \frac{d(d-1) + 2\alpha^2}{r}(V_-)_r \\
&+ \frac{(d-1)(d-2) - 2\alpha^2}{r}((V_+)_r) - (d-2)(V_-)_{rr} + 3d(V_+)_{rr} \\
&+ 2r((V_+)_rV_- + V_-(V_+)_r) + 2r(V_+)_{rrr} - 2\lambda r((V_-)_r + (V_+)_r)]u_k^2r^{d-1}dr \\
&\leq 0.
\end{aligned}$$

In the preceding calculations, we integrated by parts several times above. To justify this, $r = 0$ must be a root of u_k of sufficiently high multiplicity to compensate for the singular terms. Fortunately, spherical harmonics result from eigenvalues of the spherical Laplacian. These take the values

$$\nu_k = k^2 + (d-2)k,$$

and for each k , the eigenfunctions (and hence the spherical harmonics) are traces of harmonic polynomials of degree k . As a result, in order to give a smooth solution as guaranteed above, $u_k(0)$ must be a zero of multiplicity k , or $u_k^{(m)}(0) = 0$ for all $m = 0, 1, \dots, k$. Hence, for $k \geq \max\{0, 5-d\}$, the behavior of u is sufficient to make the calculations rigorous. See [38], Chapter 8 for a detailed description of eigenfunctions for the Laplacian on the sphere.

In the commutator expression, the parameter that must dominate is α^4 . Since V_- , V_+ are smooth, exponentially decaying functions by assumption, all terms involving V_- , V_+

and derivatives thereof are nicely bounded at 0 and exponentially decaying. Hence, for $0 \leq r \leq 1$, all of the functions above are easily controlled by $\frac{\alpha^4}{r^4}$ for α sufficiently large.

Similarly, for $r > r_*$, r_* sufficiently large, the exponential decay of V_- , V_+ and their derivatives imply that any function above is dominated $\frac{\alpha^4}{r^4}$ once α is sufficiently. In the intermediate region, using the smoothness of the potential functions we can find α^4 large enough to bound the lower order terms. In order to determine α exactly, a careful analysis must be done involving all of the extrema of V_- , V_+ and their derivatives. As these functions are uniformly bounded, there exists α_0 such that for all $\alpha \geq \alpha_0$, the operator when conjugated by the radial Mourre operator gives a negative definite system. Hence, the result holds. $\square$

Remark A.1. *Any embedded eigenvalue can be expressed purely as a finite sum of spherical harmonics with radial coefficients. Combining this with the result in Section A.1 gives a limited range for the calculations one must do in order to determine whether or not an operator has no embedded eigenvalues.*

APPENDIX B. PROOF OF THE INVERTIBILITY OF THE OPERATORS

In this section we give a full proof of Proposition 6.4. This relies on our numerically computed indexes, from Proposition 5.3. This proof generalizes to the 3d problem, establishing Proposition 5.6.

Following [13, 20], we first prove uniqueness and then existence. Before beginning the uniqueness part, we recall the following extension of the Levinson Theorem, [7, 17], from [11]:

Theorem 12 (Eastham). *For the equation*

$$y^{(n)} + a_1(x)y^{(n-1)} + \dots + a_n(x)y(x) = 0,$$

assume

$$\int_a^\infty x^{j-1}|a_j(x)|dx < \infty$$

for some $a > 0$ and $j = 1, \dots, n$. Then there exist n solutions, $y_k(x)$, such that as $x \rightarrow \infty$,

$$y_k^{(i-1)}(x) \sim \frac{x^{k-i}}{(k-i)!}, \quad 1 \leq i \leq k,$$

$$y_k^{(i-1)}(x) = o(x^{(k-i)}), \quad k+1 \leq i \leq k.$$

To prove uniqueness, let $u \in L^\infty(\mathbb{R})$ solve $\mathcal{L}u = 0$. We prove $u = 0$. The equation, $-u'' + V(x)u = 0$, satisfies the hypotheses of the above theorem, so there exist two solutions, $\rho_1(x)$ and $\rho_2(x)$, such that as $x \rightarrow +\infty$,

$$(B.1) \quad \rho_1(x) \sim 1, \quad \rho_1'(x) = o(x^{-1}),$$

$$(B.2) \quad \rho_2(x) \sim x, \quad \rho_2'(x) = o(1).$$

Due to the behaviour as $x \rightarrow +\infty$, ρ_1 and ρ_2 are linearly independent. These can then be extended to all of $\mathbb{R}$ by the classical theory of linear systems with smooth coefficients.

By the same argument, there exist $\tilde{\rho}_1$ and $\tilde{\rho}_2$ satisfying

$$(B.3) \quad \tilde{\rho}_1(x) \sim 1, \quad \tilde{\rho}'_1(x) = o(x^{-1}),$$

$$(B.4) \quad \tilde{\rho}_2(x) \sim x, \quad \tilde{\rho}'_2(x) = o(1)$$

as $x \rightarrow -\infty$. These two are also linearly independent. We will make use of these two sets of functions in what follows. An important relation between them is that, by uniqueness, $\tilde{\rho}_j(x) = \rho_j(-x)$ for $j = 1, 2$.

Given the values $u(0) = u_0$ and $u'(0) = u'_0$, we know from the existence and uniqueness of solutions to linear systems with smooth coefficients, that there exist two unique pairs of constants, $\{c_1, c_2\}$ and $\{\tilde{c}_1, \tilde{c}_2\}$ such that

$$\begin{aligned} u(x) &= c_1 \rho_1(x) + c_2 \rho_2(x) \\ &= \tilde{c}_1 \tilde{\rho}_1(x) + \tilde{c}_2 \tilde{\rho}_2(x). \end{aligned}$$

As $x \rightarrow +\infty$,

$$u(x) \sim c_1 + c_2 x.$$

Since $u(x) \in L^\infty$, we conclude $c_2 = 0$. Since it is proportional to $\rho_1(x)$, $u'(x)$ vanishes as $x \rightarrow +\infty$. An analogous argument ensures that $u'(x)$ also vanishes as $x \rightarrow -\infty$. Thus we have

$$(B.5) \quad \lim_{|x| \rightarrow \infty} u'(x) = 0.$$

Since $u \in C^2(\mathbb{R})$ and its derivative vanishes, we conclude $u' \in L^\infty(\mathbb{R})$. Furthermore, we claim $u' \in L^2(\mathbb{R})$. Multiplying the equation by u and integrating by parts,

$$\int_{-L}^L |u'|^2 dy - uu'|_L^L + \int_L^L V u^2 dy = 0.$$

Sending $L \rightarrow \infty$ proves the claim. Additionally, this shows $\langle \mathcal{L}u, u \rangle = 0$.

Let $\chi_A(x)$ be the cutoff function

$$(B.6) \quad \chi_A(x) = \begin{cases} 1, & |x| \leq A, \\ 2 \left(\frac{\log A}{\log |x|} - \frac{1}{2} \right), & A < |x| \leq A^2, \\ 0, & A^2 < |x|. \end{cases}$$

and assume $A > 1$. Let $u_A(x) = \chi_A(x)u(x)$. We prove

$$(B.7) \quad \lim_{A \rightarrow \infty} \langle \mathcal{L}u_A, u_A \rangle = \langle \mathcal{L}u, u \rangle.$$

which will be essential to the uniqueness proof. Trivially, the potential component, $\int V u_A^2$, will converge to $\int V u^2$, since V is highly localized. We now justify the convergence of the kinetic component,

$$(B.8) \quad \begin{aligned} \int |u'_A| dy &= \int \chi_A^2 |u'|^2 dy + \int |\chi'_A|^2 |u|^2 dy + 2 \int \chi_A u' \chi'_A u dy \\ &= I_1 + I_2 + I_3. \end{aligned}$$

The integral I_1 converges to $\int |u'|^2$, the desired quantity. We must show the other two vanish. First, we split up I_2 into

$$I_2 = \int |\chi'_A|^2 u^2 dy = \int_{-A^2}^{-A} |\chi'_A|^2 u^2 dy + \int_A^{A^2} |\chi'_A|^2 u^2 dy.$$

Using our explicit characterization of the cutoff function

$$\chi'_A(x) = -\frac{2 \log A}{x(\log |x|)^2},$$

$$\int_A^{A^2} |\chi'_A|^2 |u|^2 dy \leq \|u\|_{L^\infty}^2 \int_A^{A^2} \frac{4(\log A)^2}{y^2(\log y)^4} dy.$$

As $A \rightarrow +\infty$, the integral is $\sim 1/((\log(A^{-1}))^6 A)$, which vanishes as $A \rightarrow +\infty$. The integral over $(-A^2, -A)$ is treated similarly.

The other integral is

$$I_3 = 2 \int_{-A^2}^{-A} \chi_A u' \chi'_A u dy + 2 \int_A^{A^2} \chi_A u' \chi'_A u dy.$$

Again, using the explicit characterization of the cutoff function,

$$\begin{aligned} \int_A^{A^2} \chi_A u' \chi'_A u dy &\leq \|u\|_{L^\infty} \|u'\|_{L^\infty(A, A^2)} \int_A^{A^2} \frac{2 \log A}{y(\log |y|)^2} dy \\ &\leq \|u\|_{L^\infty} \|u'\|_{L^\infty(A, A^2)}. \end{aligned}$$

Since the derivative vanishes as $x \rightarrow +\infty$, (B.5), this also vanishes. The other part of I_3 is treated analogously. This proves convergence of the bilinear form.

We now specialize to either even or odd functions. By our index computations in Proposition 5.3, $\mathcal{L}_\pm^{(e/o)}$ each have index 1, except for $\mathcal{L}_-^{(o)}$ which has index 0. Without loss of generality, we assume $\mathcal{L}$ is an index 1 operator, and proceed. Let ψ be the negative eigenvector with $\|\psi\|_{L^2} = 1$ for the relevant symmetry. Let $V_A = \text{span}\{\psi, u_A\}$. We now show that $\bar{\mathcal{L}}$ restricted to this subspace is negative definite. By the index computations this will imply u_A and ψ are collinear, allowing us to conclude that $u = 0$ since u_A has the same symmetry properties to u .

Let q be any element of V_A ,

$$q = c_1 u_A + c_2 \psi.$$

Then,

$$\langle \bar{\mathcal{L}}q, q \rangle = c_1^2 \langle \bar{\mathcal{L}}u_A, u_A \rangle + 2c_1c_2 \langle \bar{\mathcal{L}}u_A, q \rangle + c_2^2 \langle \bar{\mathcal{L}}\psi, \psi \rangle.$$

We claim this is negative, which shows $\bar{\mathcal{L}}$, restricted to V_A , is negative definite. By the index computation, there is only one negative eigenvalue. Thus, $\dim V_A = 1$, and we must have $u_A = c(A)\psi$. But then

$$-\lambda c(A) = \langle u_A, \lambda\psi \rangle = \langle u_A, \mathcal{L}\psi \rangle = \langle \mathcal{L}u_A, \psi \rangle.$$

Since the right hand side vanishes as $A \rightarrow \infty$, we conclude that $c(A) = 0$; hence $u = 0$.

To prove the claim that the form is negative, it is equivalent to show

$$(B.9) \quad \langle \bar{\mathcal{L}}u_A, \psi \rangle^2 < \langle \bar{\mathcal{L}}u_A, u_A \rangle \langle \bar{\mathcal{L}}\psi, \psi \rangle.$$

As $A \rightarrow \infty$,

$$\begin{aligned} \langle \bar{\mathcal{L}}u_A, u_A \rangle &\rightarrow -\delta_0 \int e^{-|y|} |u|^2, \\ \langle \bar{\mathcal{L}}u_A, \psi \rangle &\rightarrow -\delta_0 \int e^{-|y|} u\psi \end{aligned}$$

and

$$\langle \bar{\mathcal{L}}\psi, \psi \rangle \leq \lambda < 0.$$

Thus, (B.9) holds for A sufficiently large and δ_0 sufficiently small. Indeed, given $u \neq 0$, and ψ , let

$$\delta_0 \leq \frac{1}{2} \frac{|\lambda| \int e^{-|y|} |u|^2 dy}{\int e^{-|y|} |u| |\psi| dy}.$$

Fixing this value of δ_0 , we can then find a value of A sufficiently large such that the inequality holds.

For the operator $\mathcal{L}_-^{(o)}$, the proof is simpler, as we need only observe that

$$\langle \bar{\mathcal{L}}_-^{(o)} u_A, u_A \rangle < 0,$$

contradicting the positivity of the operator. This concludes our proof of uniqueness of the solutions.

We now prove existence. Again, this follows [13, 20]. We have the two fundamental sets of solutions $\{\rho_1, \rho_2\}$ and $\{\tilde{\rho}_1, \tilde{\rho}_2\}$. ρ_1 and $\tilde{\rho}_1$ are asymptotically constant at ∞ and $-\infty$, respectively. Note that these two must be linearly independent, for if they were collinear, we would have a solution in L^∞ , solving $\mathcal{L}u = 0$. Hence,

$$\begin{aligned} |\rho_1(x)| &\leq K|x| \quad \text{as } x \rightarrow -\infty, \\ |\tilde{\rho}_1(x)| &\leq K|x| \quad \text{as } x \rightarrow +\infty \end{aligned}$$

for some constant K . We construct a Green's function from these two to get the solution

$$u(x) = \tilde{\rho}_1(x) \int_x^\infty \frac{\rho_1(s)f(s)}{W(s)} ds + \rho_1(x) \int_{-\infty}^x \frac{\tilde{\rho}_1(s)f(s)}{W(s)} ds,$$

where $W = \tilde{\rho}_1 \rho_1' - \tilde{\rho}_1' \rho_1$ is the Wronskian. The integrals converge and have the appropriate decay due to the properties of ρ_1 and $\tilde{\rho}_1$, and our assumption that f is highly localized.

Finally, if f is even, consider

$$\begin{aligned} u(-x) &= \tilde{\rho}_1(-x) \int_{-x}^{\infty} \frac{\rho_1(s)f(s)}{W(s)} ds + \rho_1(-x) \int_{-\infty}^{-x} \frac{\tilde{\rho}_1(s)f(s)}{W(s)} ds \\ &= \tilde{\rho}_1(-x) \int_{-\infty}^x \frac{\rho_1(-s)f(s)}{W(-s)} ds + \rho_1(-x) \int_x^{\infty} \frac{\tilde{\rho}_1(-s)f(s)}{W(-s)} ds \\ &= \rho_1(x) \int_{-\infty}^x \frac{\tilde{\rho}_1(s)f(s)}{W(s)} ds + \tilde{\rho}_1(x) \int_x^{\infty} \frac{\rho_1(s)f(s)}{W(s)} ds. \end{aligned}$$

Thus $u(x) = u(-x)$. An analogous proof holds for f odd.

APPENDIX C. NUMERICAL METHODS

The software tools we use in our computations are the MATLAB and Fortran 90/95 implementations of an adaptive nonlinear collocation algorithm discussed in [26–28]. Though they are quite similar, we found the Fortran implementation to be faster and more robust for solving the 3d problems which require us to compute the ground state. For the 1d problems, where we have an explicit formula, the MATLAB algorithm, `bvp4c` sufficed. We use these tools to solve for the soliton, compute the index functions, and solve the relevant boundary value problems and associated inner products.

The codes used to perform these computations are available at http://www.math.toronto.edu/simpson/files/spec_prop_code.tgz.

C.1. Singularities at the Origin. A useful feature of this algorithm is that it can handle boundary value problems of the form

$$\frac{d}{dr} \mathbf{y} = \frac{1}{r} S \mathbf{y} + \mathbf{f}(r, \mathbf{y}),$$

where S is some constant coefficient matrix. The r^{-1} singularity naturally appears in the 3D problems due to the Laplacian. The higher harmonics introduce a r^{-2} singularity which can be addressed by a change of variables. Let $\mathcal{L}^{(k)}$ denote one of the operators applied to the k -th harmonic,

$$\mathcal{L}^{(k)} = -\frac{d^2}{dr^2} - \frac{d-1}{r} \frac{d}{dr} + \mathcal{V} + \frac{k(k+d-2)}{r^2}.$$

Then if W solves $\mathcal{L}^{(k)} W = f$, where f could be zero and W is non singular at the origin, then

$$\lim_{r \rightarrow 0} r^{-k} W(r)$$

is a nonzero constant. This motivates the change of variable $W(r) = r^k \widetilde{W}(r)$. In terms of $\widetilde{W}(r)$, the equation becomes

$$r^k \widetilde{\mathcal{L}}^{(k)} \widetilde{W} = f,$$

where

$$\widetilde{\mathcal{L}}^{(k)} = -\frac{d^2}{dr^2} - \frac{d-1+2k}{r} \frac{d}{dr} + \mathcal{V}$$

and $\widetilde{W}$ satisfies the condition $\widetilde{W}'(0) = 0$. We compute with $\widetilde{\mathcal{L}}^{(k)}$ to get $\widetilde{W}$ and then multiply by r^k . When we compute indexes for these higher harmonic operators, the other initial condition becomes $\widetilde{W}(0) = 1$.

C.2. Artificial Boundary Conditions. Another subtlety of the computations is the far field boundary conditions. The soliton, R , vanishes as $r \rightarrow +\infty$, but we only compute out to some finite value, $r_{\max}$. For simplicity, we use the notation r and $r_{\max}$ for both 1D and 3D. To accommodate this, we introduce an artificial boundary condition at $r_{\max}$, and then check *a posteriori*, that it is consistent. Thus, we must do some asymptotic analysis.

We seek an asymptotic expansion for R , using (2.1)

$$(-\Delta + \lambda - f(R))R = 0.$$

As $r \rightarrow \infty$, we look for an expansion of the form

$$(C.1) \quad e^{-\sqrt{\lambda}r} r^\gamma \sum_{n=0}^{\infty} c_n r^{-n}.$$

To extract the leading order behavior, we wish to find γ . To this end, we have

$$\begin{aligned} \frac{\partial}{\partial r}(e^{-\sqrt{\lambda}r} r^\gamma) &= -\sqrt{\lambda} e^{-\sqrt{\lambda}r} r^\gamma + \gamma e^{-\sqrt{\lambda}r} r^{\gamma-1}, \\ \frac{\partial^2}{\partial^2 r}(e^{-\sqrt{\lambda}r} r^\gamma) &= \lambda e^{-\sqrt{\lambda}r} r^\gamma - 2\lambda\gamma e^{-\sqrt{\lambda}r} r^{\gamma-1} + \gamma(\gamma-1) e^{-\sqrt{\lambda}r} r^{\gamma-2}. \end{aligned}$$

Plugging (C.1) into (2.1), we see

$$[(-\lambda + \lambda) + \frac{(2\sqrt{\lambda}\gamma + \sqrt{\lambda}(d-1))}{r} + O(r^{-2})] = 0.$$

Hence, $\gamma = -(d-1)/2$ and the leading order behavior is

$$(C.2) \quad r^{-\frac{d-1}{2}} e^{-\sqrt{\lambda}r}.$$

Therefore, as $r \rightarrow \infty$,

$$(C.3) \quad R(r) \approx \begin{cases} R_\star e^{-\sqrt{\lambda}r} & d = 1, \\ R_\star \frac{1}{r} e^{-\sqrt{\lambda}r} & d = 3. \end{cases}$$




FIGURE 13. The numerically computed ground state for the different problems. Computed on the indicated domain, with a tolerance of 10^{-13} while assessing the indexes of the operators restricted to even functions for $d = 1$ and the zeroth harmonic for $d = 3$. The computed solitons are monotonic and decay at the anticipated rate.

From this, we develop the Robin boundary condition,

$$\lim_{r \rightarrow \infty} \frac{R(r)}{R'(r)} \rightarrow \begin{cases} -\frac{1}{\sqrt{\lambda}} & d = 1, \\ -\frac{r}{1 + \sqrt{\lambda}r} & d = 3, \end{cases}$$

which we formulate as

$$(C.4a) \quad R(r_{\max}) + \frac{1}{\sqrt{\lambda}} R'(r_{\max}) = 0 \quad \text{for } d = 1$$

and

$$(C.4b) \quad R(r_{\max}) + \frac{r_{\max}}{1 + \sqrt{\lambda}r_{\max}} R'(r_{\max}) = 0 \quad \text{for } d = 3$$

assuming we have taken $r_{\max}$ sufficiently large. For our computations, aside from noting that the solver algorithm ends without errors, we have two *a posteriori* checks available. The first is to verify that we have, in fact, computed the ground state. Plotting the computed R on both a linear and a log scale in Figure 13 we verify that R is a hump shaped monotonically decaying function. It also has the anticipated $r^{-1}e^{-r}$ decay rate.

The second thing that can be checked is that the numerically computed R satisfies, asymptotically, the artificial boundary condition (C.4b). To do this, we plot $R(r) + \frac{r}{1 + \sqrt{\lambda}r} R'(r)$ and observe that it vanishes as $r \rightarrow \infty$. As can be seen in Figure 14, the mismatch in the artificial boundary condition is small and monotonically decaying in r .




FIGURE 14. As $r \rightarrow \infty$, R asymptotically satisfies $-\Delta R + \lambda R \sim 0$. The figures indicate that the relevant domain, which is different for different problems, is sufficiently large that artificial boundary conditions are good approximations. From the same computation as in Figure 13.

To compute the off axis eigenstates $\vec{\phi}$ for the 3D cubic problem with eigenvalue $\sigma > 0$, we rely on the relationship

$$L_- L_+ \phi_1 = -\sigma^2 \phi_1.$$

Asymptotically, this is the free equation

$$(-\Delta + 1)^2 \phi_1 = -\sigma^2 \phi_1.$$

Seeking a radially symmetric solution and using similar expansion techniques as in the case of the soliton, we will find that

$$(C.5) \quad \phi_1(r) \approx c_1 r^{-(d-1)/2} e^{-r\sqrt{1+i\sigma}} + c_2 r^{-(d-1)/2} e^{-r\sqrt{1-i\sigma}}$$

as $r \rightarrow \infty$.

Let

$$(C.6) \quad \theta \equiv \frac{1}{2} \arctan(\sigma),$$

$$(C.7) \quad \rho \equiv (1 + \sigma^2)^{1/4}.$$

In 1D, we can construct an artificial boundary condition that, as $r \rightarrow r_{\max}$,

$$(C.8) \quad \phi_1' + \rho \cos(\theta) \phi_1 + \rho \sin(\theta) \phi_2 = 0,$$

$$(C.9) \quad \phi_2' + \rho \cos(\theta) \phi_2 - \rho \sin(\theta) \phi_1 = 0.$$




FIGURE 15. The numerically computed ground state and the off axis unstable mode of 3d cubic NLS. Computed on the indicated domain, with a tolerance of 10^{-12} .

Analogously, in 3D, as $r \rightarrow r_{\max}$,

$$(C.10) \quad \phi_1' + \rho \cos(\theta) \phi_1 + \frac{1}{r} \phi_1 + \rho \sin(\theta) \phi_2 = 0,$$

$$(C.11) \quad \phi_2' + \rho \cos(\theta) \phi_2 + \frac{1}{r} \phi_2 - \rho \sin(\theta) \phi_1 = 0.$$

As in the case of the soliton, we can verify that the solutions have the appropriate shape, decay as expected, and satisfy the artificial boundary conditions. The shape and decay are plotted in Figure 15. The functions rapidly reach machine precision. If we zoom in on the unstable modes, as in Figure 16, we can see the periodic structure. However, as is suggested by these figures, once ϕ_j 's are sufficiently small, $\lesssim O(10^{-12})$, this fine structure degrades. Fortunately, this numerical error is sufficiently small as to not impact our computations. The artificial boundary condition plot appears in Figure 17.

Another place where we use artificial boundary conditions is in solving the various boundary value problems for $U_j^{(k)}$ or $Z_j^{(k)}$. The rapid decay of the soliton leads to the functions satisfying the free equation, for $d > 1$,

$$-\Delta q + k(k + d - 2)/r^{d-2+k} q = 0,$$

where q is one of these functions. In this region, the function must asymptotically be like (5.7b). Of course, we work with the variables $\tilde{U}_j^{(k)}$ and $\tilde{Z}_j^{(k)}$. Since these vanish as $r \rightarrow \infty$, they asymptotically behave as

$$q \propto r^{2-d-2k}.$$




FIGURE 16. Zooming in on the components of the numerically computed unstable mode, $\vec{\phi}$, we see the periodic structure of the solution.

Thus, we have the artificial boundary conditions, valid for all harmonics and $d > 2$,

$$(C.12a) \quad \tilde{U}_j^{(k)}(r_{\max}) + \frac{r}{d-2+2k} \frac{d}{dr} \tilde{U}_j^{(k)}(r_{\max}) = 0,$$

$$(C.12b) \quad \tilde{Z}_j^{(k)}(r_{\max}) + \frac{r}{d-2+2k} \frac{d}{dr} \tilde{Z}_j^{(k)}(r_{\max}) = 0.$$

Figure 17 shows that these artificial boundary conditions are asymptotically satisfied.

For our $d = 1$ computations, we have the artificial boundary conditions

$$(C.13) \quad \frac{d}{dx} U_j^{(e/o)}(x_{\max}) = 0, \quad \frac{d}{dx} Z_j^{(e/o)}(x_{\max}) = 0.$$

We can similarly check that the inner products are asymptotically constant and that our computed functions asymptotically satisfy the artificial boundary conditions. See Figures 18 and C.2.




FIGURE 17. As $r \rightarrow \infty$, our functions asymptotically satisfy the free, linear equations for the different parts of the 3d cubic problem. The figures indicate that the relevant domain, which is different for different problems, is sufficiently large that artificial boundary conditions are good approximations.




FIGURE 18. As $r \rightarrow \infty$, our functions asymptotically satisfy the free equations for the various 1d supercritical problems.




FIGURE 19. As $r \rightarrow \infty$, our functions asymptotically satisfy the free equations for the 1d critical problem with various orthogonality conditions.

The last place we make use of artificial boundary conditions is in solving $L_+\beta = -x^2R$ for the critical problem in Section 6.2.3. Using the same procedure as above, one will find that $\beta \propto x^3e^{-x}$ form which the artificial boundary condition

$$(C.14) \quad \beta'(r) + \left(1 - \frac{3}{r}\right)\beta(r) = 0$$

can be constructed.

C.3. Computation of the Indexes. In computing the indexes of operators $\tilde{\mathcal{L}}_{\pm}^{(k)}$ (from which we recover the indexes of $\mathcal{L}_{\pm}^{(k)}$), we simultaneously solve the mixed boundary value

problem/initial value problems

$$(C.15) \quad -\Delta R + \lambda R - g(|R|^2)R = 0, \quad R'(0) = 0, \quad (C.4a) \text{ or } (C.4b),$$

$$(C.16) \quad \tilde{\mathcal{L}}_+^{(k)} \tilde{U}^{(k)} = 0, \quad \tilde{U}^{(k)}(0) = 1, \quad \frac{d}{dr} \tilde{U}^{(k)}(0) = 0,$$

$$(C.17) \quad \tilde{\mathcal{L}}_-^{(k)} \tilde{Z}^{(k)} = 0, \quad \tilde{Z}^{(k)}(0) = 1, \quad \frac{d}{dr} \tilde{Z}^{(k)}(0) = 0.$$

For the 3d cubic equation, this is the complete set of equations; $\lambda = 1$ and $f(s) = s$. The analogous computations are made in $d = 1$. In addition to verifying that the soliton was adequately computed, we can check, *a posteriori*, that the index functions, $U^{(k)}$ and $Z^{(k)}$ asymptotically satisfy the free equation. This was shown in the index figures of Sections 5.2 and 6.1, where we checked the constants.

C.4. Computation of the Inner Products. As in the computation of the indexes, we similarly solve mixed boundary value/initial value problem for R , the $U_j^{(\alpha)}$ and $Z_j^{(\alpha)}$, and the inner products

In computing the inner products, we introduce the dependent variables $\kappa_j^{(\alpha)}(r)$ and $\gamma_j^{(\alpha)}(r)$, where

$$(C.18) \quad \frac{d}{dr} \kappa_j^{(\alpha)}(r) = \mathcal{L}_+^{(\alpha)} U_{\ell_1}^{(\alpha)} U_{\ell_2}^{(\alpha)} r^{d-1}, \quad \kappa_j^{(\alpha)}(0) = 0,$$


$$(C.19) \quad \frac{d}{dr} \gamma_j^{(\alpha)}(r) = \mathcal{L}_-^{(\alpha)} Z_{\ell_1}^{(\alpha)} Z_{\ell_2}^{(\alpha)} r^{d-1}, \quad \gamma_j^{(\alpha)}(0) = 0$$

for ℓ_1 and ℓ_2 the appropriate indexes. Clearly,


$$\lim_{r \rightarrow \infty} \kappa_j^{(\alpha)}(r) = K_j^{(\alpha)},$$

$$\lim_{r \rightarrow \infty} \gamma_j^{(\alpha)}(r) = J_j^{(\alpha)}.$$


We approximate the inner products by computing to $r_{\max}$. As demonstrated in Figures 20, 21, 22 these converge rapidly and are essentially constant in the region of the free equation. This is entirely consistent with the exponential decay of the soliton and functions related to it, such as its derivative.



(a) Inner Products for $\mathcal{L}_+^{(0)}$ functions for 3D Cubic



(b) Inner Products for $\mathcal{L}_-^{(0)}$ functions for 3D Cubic



(c) Inner Products for $\mathcal{L}_+^{(1)}$ functions for 3D Cubic

FIGURE 20. Inner products for the 3d cubic equation.




FIGURE 21. Inner products for the 1d supercritical problems




FIGURE 22. Inner products for the 1d critical problem with various orthogonality conditions.

REFERENCES

- [1] S. Agmon. Spectral properties for Schrödinger operators and scattering theory. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 2:151–218, 1975.
- [2] H. Berestycki and P.L. Lion. Nonlinear scalar field equations, I: Existence of a ground state. *Arch. Rational. Mech. anal.*, 82(4):313–345, 1983.
- [3] V. S. Buslaev and G. S. Perel'man. On the stability of solitary waves for nonlinear Schrödinger equations. In *Nonlinear evolution equations*, volume 164 of *Amer. Math. Soc. Transl. Ser. 2*, pages 75–98. Amer. Math. Soc., Providence, RI, 1995.
- [4] Vladimir S. Buslaev and Catherine Sulem. On asymptotic stability of solitary waves for nonlinear Schrödinger equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 20(3):419–475, 2003.
- [5] T. Cazenave. *Semilinear Schrödinger Equations*, volume 10 of *Courant Lecture Notes in Mathematics*. American Mathematical Society, 2003.
- [6] Shu-Ming Chang, Stephen Gustafson, Kenji Nakanishi, and Tai-Peng Tsai. Spectra of linearized operators for NLS solitary waves. *SIAM J. Math. Anal.*, 39(4):1070–1111, 2007/08.
- [7] Earl A. Coddington and Norman Levinson. *Theory of Ordinary Differential Equations*. Krieger, 1984.

- [8] Scipio Cuccagna. On asymptotic stability of ground states of NLS. *Rev. Math. Phys.*, 15(8):877–903, 2003.
- [9] Scipio Cuccagna and Dmitry Pelinovsky. Bifurcations from the endpoints of the essential spectrum in the linearized nonlinear Schrödinger problem. *J. Math. Phys.*, 46(5):053520, 15, 2005.
- [10] L. Demanet and W. Schlag. Numerical verification of a gap condition for a linearized nonlinear Schrödinger equation. *Nonlinearity*, pages 829–852, 2006.
- [11] M.S.P. Eastham. *The Asymptotic Solution of Linear Differential Systems: Application of the Levinson Theorem*. Oxford University Press, 1989.
- [12] B. Erdogan and W. Schlag. Dispersive estimates for Schrödinger operators in the presence of a resonance and/or eigenvalue at zero energy in dimension three. II. *J. Anal. Math.*, 99:199–248, 2006.
- [13] G. Fibich, F. Merle, and P. Raphaël. Proof of a Spectral Property related to the singular formation for the L^2 critical nonlinear Schrödinger equation. *Physica D*, 220:1–13, 2006.
- [14] M. Grillakis, J. Shatah, and W. Strauss. Stability theory of solitary waves in the presence of symmetry. ii. *J. Funct. Anal.*, 94(2):308–348, 1990.
- [15] P. Hislop and I.M. Sigal. *Introduction to Spectral Theory With Applications to Schrödinger Operators*, volume 113 of *Applied Mathematical Sciences*. Springer-Verlag, 1996.
- [16] J. Krieger and W. Schlag. Stable manifolds for all monic supercritical focusing nonlinear Schrödinger equations in one dimension. *J. Amer. Math. Soc.*, 19(4):815–920, 2006.
- [17] Norman Levinson. The asymptotic nature of solutions of linear systems of differential equations. *Duke Mathematical Journal*, 15:111–126, 1948.
- [18] J. Marzuola. Dispersive estimates using scattering theory for matrix Hamiltonian equations. Preprint arXiv:0906.0351, 2009.
- [19] K. McCleod. Uniqueness of Positive Radial Solutions of $\Delta u + f(u) = 0$ in $\mathbb{R}^n$, II. *Transactions of the American Mathematical Society*, 339(2):495–505, 1993.
- [20] F. Merle and P. Raphael. The blow-up dynamic and upper bound on the blow-up rate for critical nonlinear Schrödinger equation. *Annals of Mathematics*, pages 157–222, 2005.
- [21] Galina Perelman. *Personal communication to Wilhelm Schlag*.
- [22] Galina Perelman. On the formation of singularities in solutions of the critical nonlinear Schrödinger equation. *Ann. Henri Poincaré*, 2(4):605–673, 2001.
- [23] M. Reed and B. Simon. *Methods of Modern Mathematical Physics IV. Analysis of Operators*. Academic Press, 1978.
- [24] I. Rodnianski, W. Schlag, and A. Soffer. Asymptotic stability of N -soliton states of NLS. Preprint, 2003.
- [25] W. Schlag. Stable manifolds for an orbitally unstable NLS. To appear in *Annals of Math*, 2004.
- [26] L.F. Shampine. Singular boundary value problems for ODEs. *Applied Mathematics and Computation*, 138(1):99–112, 2003.
- [27] L.F. Shampine, I. Gladwell, and S. Thompson. *Solving ODEs with MATLAB*. Cambridge University Press, 2003.
- [28] L.F. Shampine, P.H. Muir, and H. Xu. A User-Friendly Fortran BVP Solver. *JNAIAM*, 1(2):201–217, 2006.
- [29] J. Shatah. Stable Standing Waves of Nonlinear Klein-Gordon Equations. *Communications in Mathematical Physics*, 91:313–327, 1983.
- [30] J. Shatah. Unstable Ground State of Nonlinear Klein-Gordon Equations. *Transactions of the American Mathematical Society*, 290(2):701–710, 1985.
- [31] J. Shatah and W. Strauss. Instability of Nonlinear Bound States. *Communications in Mathematical Physics*, 100:173–190, 1985.
- [32] G. Simpson and M.I. Weinstein. Asymptotic stability of ascending solitary magma waves. *SIAM J. Math. Anal.*, 40:1337–1391, 2008.

- [33] E. Stein. *Harmonic Analysis: real-variable methods, orthogonality, and oscillatory integrals*, volume 43 of *Princeton Mathematical Series*. Princeton University Press, 1993.
- [34] C. Sulem and P. Sulem. *The Nonlinear Schrödinger Equation. Self-focusing and wave-collapse*, volume 39 of *Applied Mathematical Sciences*. Springer-Verlag, 1999.
- [35] C. Sulem and P.L. Sulem. *The Nonlinear Schrödinger Equation: Self-Focusing and Wave Collapse*. Springer, 1999.
- [36] Terence Tao. A global compact attractor for high-dimensional defocusing non-linear Schrödinger equations with potential. *Dyn. Partial Differ. Equ.*, 5(2):101–116, 2008.
- [37] D. Tataru. Microlocal Analysis. Unpublished Lecture Notes, 2003.
- [38] M. Taylor. *Partial Differential Equations II*, volume 116 of *Classics in Mathematics*. Springer-Verlag, 1997.
- [39] M. I. Weinstein. Modulational stability of ground states of nonlinear Schrödinger equations. *SIAM J. Math. Anal.*, 16:472–491, 1985.
- [40] M. I. Weinstein. Lyapunov stability of ground states of nonlinear dispersive evolution equations. *Comm. Pure Appl. Math.*, 39:472–491, 1986.

E-mail address: jm3058@columbia.edu

APPLIED MATHEMATICS DEPARTMENT, COLUMBIA UNIVERSITY, 200 S. W. MUDD, 500 W. 120TH ST., NEW YORK CITY, NY 10027, USA

E-mail address: simpson@math.toronto.edu

MATHEMATICS DEPARTMENT, UNIVERSITY OF TORONTO, TORONTO, ONTARIO, CANADA